VOLUME 37 • AUTHOR INDEX

Aaltonen LA *see* Roth S *et al*
see Wheeler JMD *et al*
Abe S *see* Usami S-I *et al*
Abe S *et al*. Prevalent connexin 26 gene (*GJB2*) mutations in Japanese, **41**
Abidi F *see* Cabezas DA *et al*
Abramowicz MJ *et al*. Fetal bowel hyperechogenicity may indicate mild atypical cystic fibrosis: a case associated with a complex *CFTR* allele: *letter*, **e15**
Abu-Amero SN *see* Wakeling EL *et al*
Adès L *see* Reardon W *et al*
Adinolfi M, Cirigliano V. Detection of fetal cells in transcervical samples using X22 marker: *letter*, **e1**
Affara N *see* Blanco P *et al*
Affara NA *see* Sargent CA *et al*
see Woodward ER *et al*
Aftimos S *see* Fryns J-P and Aftimos S
see Robertson SP *et al*
Agnarsson BA *see* Sigbjörnsdottir BI *et al*
Ain MC *see* Wright MJ *et al*
Ajzenberg C *see* Ion A *et al*
Akaiwa M *see* Gao P-S *et al*
Akçören Z *see* Dinçer P *et al*
Akita J *see* Usami S-I *et al*
Aktan-Collan K *et al*. Evaluation of a counselling protocol for predictive genetic testing for hereditary non-polyposis colorectal cancer, **108**
Albrecht B *see* Wieczorek D *et al*
Al-Dabbagh M *see* Meeks M *et al*
Aldred MA *et al*. Germline mosaicism for a *GNAS1* mutation and Albright hereditary osteodystrophy: *letter*, **e35**
Al-Hathal MM *see* Amir IM *et al*
Allard C *see* Rivard SR *et al*
Allcock R *see* Thomson JR *et al*
Alloisio N *see* Morlé L *et al*
Al-Mahdawi SAH *see* Pook MA *et al*
Al-Tawil KI *see* Amir IM *et al*
Amiel J *see* Cormier-Daire V *et al*
see Sanlaville D *et al*
Amir IM *et al*. Deletion (2)(p14p15) in a child with severe neurodevelopmental delay: *letter*, **e21**
Anglim C *see* Gray J *et al*
Antiñolo G *see* Bessant DAR *et al*
see Borrego S *et al*
see Marcos I *et al*
Anton-Culver H *see* Seltmann M *et al*
Antony JA *see* Ellaway CJ *et al*
Anvret M *see* Xiang F *et al*
Apak MY *see* Nuytinck L *et al*
Appleton R *see* Pook MA *et al*
Arab SB *see* Masmoudi S *et al*
Arden-Jones A *see* Forrest MS *et al*
Arena JF *see* Cabezas DA *et al*
Arinobu Y *see* Gao P-S *et al*
Armour CM *et al*. Clinical and radiological assessment of a family with mild brachydactyly type A1: the usefulness of metacarpophalangeal profiles, **292**
Armstrong-Buisseret L *see* Cross SJ *et al*
Arnold N *see* Weimer J *et al*
Ashton M *see* Marsh AJ *et al*
Assouline B *see* Bahuau M *et al*
Astrinidis A *et al*. Mutational analysis of the tuberous sclerosis gene *TSC2* in patients with pulmonary lymphangioleiomyomatosis, **55**
Astuti D *see* Woodward ER *et al*
Astuto LM *see* Pieke-Dahl S *et al*
Au K-S *see* Astrinidis A *et al*
Aubin G *see* Rivard SR *et al*
Aubry MC *see* Sanlaville D *et al*
Audit *see* Hodgson SV *et al*
Audrezet M-P *see* Chen J-M *et al*
Ausems MGEM *see* Terhal PA *et al*
Ayadi H *see* Masmoudi S *et al*

Baboonian C *see* D'Cruz LG *et al*
Badzioch M *et al*. Suggestive evidence for a site specific prostate cancer gene on chromosome 1p36: *letter*, **947**
Bagci G *see* Kotzot D *et al*
Bagshaw RJ *see* Aldred MA *et al*
Bahuau M *et al*. Tandem duplication within the neurofibromatosis type 1 gene (*NF1*) and reciprocal t(15;16)(q26.3;q12.1) translocation in familial association of NF1 with intestinal neuronal dysplasia type B (IND B): *letter*, **146**
Bailey MES *see* Xiang F *et al*
Baird C *see* Whiteford ML *et al*
Bakker L *see* Eussen BHJ *et al*
Balestri P *see* Eussen BHJ *et al*
Balmer D *see* Röthlisberger B *et al*
Baraitser M *see* Reardon W *et al*
Baralle D *et al*. Benign familial infantile convulsions: report of a UK family and confirmation of genetic heterogeneity: *letter*, **e31**
Bar-Am I *see* Lev D *et al*
Bardoni B *see* Salat U *et al*
Barkardottir RB *see* Sigbjörnsdottir BI *et al*
Barnier A *see* Vuillaumier-Barrot S *et al*
Barnoux M *see* Ion A *et al*
Barratt A *see* Meiser B *et al*
Barresi R *et al*. Disruption of heart sarcoglycan complex and severe cardiomyopathy caused by β sarcoglycan mutations, **102**
Barrett TG *et al*. The mitochondrial genome in Wolfram syndrome: *letter*, **463**
Bartalini G *see* Eussen BHJ *et al*
Barton G *see* Gray J *et al*
Basaran S *see* Brecevic L *et al*
see Kotzot D *et al*
Baser ME *see* Evans DGR *et al*
see Evans DGR *et al*
Baser ME *et al*. Clinical and molecular correlates of somatic mosaicism in neurofibromatosis 2: *letter*, **542**
Basha NJ *see* Koorey D *et al*
Bashir R *see* Lindsey JC *et al*
Bassères DS *see* Vasconcellos JPC *et al*
Baumer A *see* Kotzot D *et al*
Baumstark A *see* Jakubowski L *et al*
Beach R *see* Baralle D *et al*
Beales P *see* Warburg M *et al*
Beckmann JS *see* Dinçer P *et al*
Bednarz A *see* Barrett TG *et al*
Beensen V *see* Heller A *et al*
Beesley CE *et al*. Mutational analysis of Sanfilippo syndrome type A (MPS IIIA): identification of 13 novel mutations: *letter*, **704**
Bell R *et al*. Reduction of the genetic interval for lymphoedema-distichiasis to below 2 Mb: *letter*, **725**
Bellus GA *see* Wright MJ *et al*
Beneyto M *see* Sánchez F *et al*
Bennetts B *see* Henderson S *et al*
Beresford RG *et al*. Absence of fragile X syndrome in Nova Scotia: *letter*, 77
Berg JN *et al*. A family with hereditary port wine stain: *letter*, **e12**
Berry V *see* Francis PJ *et al*
Besley GTN *see* Sargent CA *et al*
Bessant DAR *et al*. Importance of the autosomal recessive retinitis pigmentosa locus on 1q31-q32.1 (RP12) and mutation analysis of the candidate gene *RGS16* (*RGS-r*): *letter*, **384**
Besson R *see* Manouvrier-Hanu S *et al*
Bhattacharya SS *see* Bessant DAR *et al*
see Francis PJ *et al*
Bignon Y-J *see* Vaurs-Barrière C *et al*
Binkert F *see* Kotzot D *et al*
Birch P *see* Szudek *et al*
Bird AC *see* Bessant DAR *et al*
see Webster AR *et al*
Bishop DT *see* Badzioch M *et al*
see Forrest MS *et al*
Bitner-Glindzicz M *see* Rickard S *et al*
Black PMC *see* Zhou X-P *et al*
Blanchard S *see* Masmoudi S *et al*
Blanck C *see* Knoblauch H *et al*
Blanck C *et al*. Three novel *SALL1* mutations extend the mutational spectrum in Townes-Brocks syndrome: *letter*, **303**
Blanco P *et al*. Divergent outcomes of intrachromosomal recombination on the human Y chromosome: male infertility and recurrent polymorphism, **752**
Blanton S *see* Kelly TE *et al*
Blaschak CJ *see* Marcos I *et al*
Blau H *see* Meeks M *et al*
Blough RI *see* Petrij F *et al*
Bocian E *see* Stankiewicz P *et al*
Bodmer WF *see* Wheeler JMD *et al*
Bolton PF *see* de Vries PJ and Bolton PF
Bolt J *see* de Lange RPJ *et al*
Bonduelle M *see* Janssens K *et al*
Boner AL *see* Patuzzo C *et al*
Bonne G *see* Forissier J-F *et al*
Bonnet V *see* Cudry S *et al*
Bonnet D *see* Cross SJ *et al*
Book reviews
Cohen M Jr, MacLean RE, *editors*. Craniosynostosis. Diagnosis, evaluation and management, **727**
Gerson SL, Keagle MB, *editors*. The Principles of Clinical Cytogenetics, **399**

Warburg M. Eye and Face in Syndromes—The Clinical Examination of Eyes and their Surroundings. Video, **399**
Boot-Handford RP *see* Gregory CA *et al*
Bordin S *see* Vasconcellos JPC *et al*
Borrego S *see* Marcos I *et al*
Borrego S *et al*. *RET* genotypes comprising specific haplotypes of polymorphic variants predispose to isolated Hirschsprung disease, **572**
Bošák V *see* Zatková A *et al*
Bottani A *see* Blanck C *et al*
Boulandet EG *see* Abramowicz MJ *et al*
Bousono C *see* Özkinay F *et al*
Bouzouita J-E *see* Masmoudi S *et al*
Bowman MC *see* Wang J *et al*
Boxer M *see* Rickard S *et al*
Boye E *see* Plant KE *et al*
Boyne J *et al*. Many ΔF508 heterozygote neonates with transient hypertrypsinaemia have a second, mild *CFTR* mutation: *letter*, **543**
Bozon D *see* Cudry S *et al*
Bozon M *see* Morlé L *et al*
Bradley T *see* Westman J *et al*
Bradshaw N *see* Campbell H *et al*
Braga S *see* Sobetzko D *et al*
Brain K *see* Gray J *et al*
Brain K *et al*. Why do women attend familial breast cancer clinics?, **197**
Brancazio LR *see* Ndah BV *et al*
Branston L *see* Gray J *et al*
Brayne C *see* Narain Y *et al*
Brecevic L *see* Kotzot D *et al*
Brecevic L *et al*. Tandem triplication of chromosome 13q14 with inverted interstitial segment in a 4 year old girl: *letter*, **964**
Brems H *see* López-Correa C *et al*
Breuning MH *see* Petrij F *et al*
Briard M-L *see* Cormier-Daire V *et al*
Briault S *see* Forissier J-F *et al*
Bricarelli FD *see* Perfumo C *et al*
Brice G *see* Bell R *et al*
Briggs MD *see* Newman B *et al*
Brik R *see* Janssens K *et al*
Brink P *see* Moolman-Smook J *et al*
Brodeur GM *see* Frühwald MC *et al*
Brondum-Nielsen K *see* Bugge M *et al*
Brook JD *see* Cross SJ *et al*
Browne CE *et al*. Duplication of medial 15q confirmed by FISH: *letter*, **e10**
Browne CE *see* Marsh AJ *et al*
Brugnoni R *see* Barresi R *et al*
Brunner HG *see* Ion A *et al*
Brusgaard K *see* Thiart R *et al*
Bruttini M *et al*. Mosaicism in Alport syndrome and genetic counselling: *letter*, **717**
Bruun-Petersen G *see* Bugge M *et al*
Bruzzone D *see* Montera M *et al*
Buervenich S *see* Xiang F *et al*
Bugge M *et al*. Disease associated balanced chromosome rearrangements: a resource for large scale genotype-phenotype delineation in man, **858**
Bulman DE *see* Armour CM *et al*
Burfeind P *see* Blanck C *et al*
Burge D *see* Marsh AJ *et al*
Burghes AHM *see* Coovert DD *et al*
Burn J *see* Lynch SA *et al*
Burnand K *see* Bell R *et al*
Burton KA *et al*. Autosomal translocation associated with premature ovarian failure: *letter*, **e2**
Burton M *see* Scheffer H *et al*
Burwinkel B *et al*. Phosphorylase kinase deficient liver glycogenosis: progression to cirrhosis in infancy associated with *PHKG2* mutations (H144Y and L225R): *letter*, **376**
Bushby KMD *see* Lindsey JC *et al*
Butow P *see* Meiser B *et al*
Buys CHCM *see* Scheffer H *et al*
Byatt S-A *see* Sidwell RU *et al*
Byron K *see* Halliday JL *et al*

Cabezas DA *et al*. A new X linked mental retardation (XLMR) syndrome with short stature, small testes, muscle wasting, and tremor localises to Xq24-q25, **663**
Cairns N *see* Narain Y *et al*
Cali A *see* Perfumo C *et al*
Caligo M *see* Hodgson SV *et al*
Callen DF *see* Bell R *et al*
Calvo C *see* Özkinay F *et al*
Cameron J *see* Ellis D *et al*
Campbell H *et al*. Evidence based medicine in practice: lessons from a Scottish clinical genetics project, **684**
Campbell A *see* Slavotinek AM *et al*
Camper SA *see* Martin DM *et al*
Cappelli M *see* Humphreys L *et al*
Cardoso C *et al*. *ATR-X* mutations cause impaired nuclear location and altered DNA binding properties of the XNP/ATR-X protein, **746**
Carlin J *see* Massie J *et al*
Carrier L *see* Forissier J-F *et al*
Carsillo T *see* Astrinidis A *et al*
Carter ND *see* D'Cruz LG *et al*
see Heathcote K *et al*
Cartigny M *see* Manouvrier-Hanu S *et al*
Casals *see* Kilinç *et al*
Casson D *see* Aldred MA *et al*
Castellan C *see* Kotzot D *et al*
Catchpoole D *et al*. Mutation analysis of *H19* and *NAP1L4* (*hNAP2*) candidate genes and *IGF2* DMR2 in Beckwith-Wiedemann syndrome: *letter*, **212**
Cavani S *see* Perfumo C *et al*
Ceccherini I *see* Costa M *et al*
Çelebi JT *et al*. Identification of *PTEN* mutations in metastatic melanoma specimens, **653**
Chadwick RB *see* Desai DC *et al*
Chamberlain S *see* Pook MA *et al*
Chan LYS *see* Lam C-W *et al*
Chan WK *see* Lam FWF *et al*
Chang-Claude J *see* Hodgson SV *et al*
see Plaschke J *et al*
Chan K-Y *see* Lam C-W *et al*
Chaoui R *see* Knoblauch H *et al*
Chapman C *et al*. Genetics of club foot in Maori and Pacific people, **680**
Charbonnier B *see* Forissier J-F *et al*
Chauvet M-L *see* Cormier-Daire V *et al*
Cheehab M *see* Meeks M *et al*
Chen J-M *et al*. A CGC>CAT gene conversion-like event resulting in the R122H mutation in the cationic trypsinogen gene and its implication in the genotyping of pancreatitis: *letter*, **e36**
Mutational analysis of the human pancreatic secretory trypsin inhibitor (*PSTI*) gene in hereditary and sporadic chronic pancreatitis: *letter*, **67**
Chen JYC *see* Goh DLM *et al*
Chen X-N *see* Slavotinek AM *et al*
Cheung KC *see* Howell A *et al*
Chiarelli I *see* Kilinç *et al*
Child AH *see* Bell R *et al*
Ching Y-H *see* Cross SJ *et al*
Chi H *see* Dahia PM *et al*
Chorley W *see* Ellis D *et al*
Chrzanowska K *see* Kotzot D *et al*
see Röthlisberger B *et al*
Chu WP *see* Lam FWF *et al*
Chudoba I *see* Heller A *et al*
Chung E *see* Meeks M *et al*
Cirigliano V *see* Adinolfi M and Cirigliano V
Civitelli S *see* Montera M *et al*
Clarke A *see* Brain K *et al*
see Elwyn G *et al*
see Gray J *et al*
Claussen U *see* Heller A *et al*
Clayton-Smith J *see* Slavotinek AM *et al*
Cleaver JE *see* Savarirayan R *et al*
Cleiren E *see* Wuyts W *et al*
Clement-Jones M *see* Jones EA *et al*
Clifford SC *see* Woodward ER *et al*
Clough MV *see* Wright MJ *et al*
Coates BJ *see* Hart PS *et al*
Cobben JM *see* Blanck C *et al*
Coble C *see* Stone DL *et al*
Cockwell AE *et al*. An analphoid supernumerary marker chromosome derived from chromosome 3 ascertained in a fetus with multiple malformations: *letter*, **807**
Cohen D *see* Gray J *et al*
Cohen BB *see* Young IE *et al*
Cohn DH *see* Mortier GR *et al*
Colleaux L *see* Cardoso C *et al*
Collet L *see* Morlé L *et al*
Colley A *see* Meiser B *et al*
Collin JRO *see* Bell R *et al*
Colombo D *see* Villa N *et al*
Commer T *see* Plaschke J *et al*
Compe E *see* Cardoso C *et al*
Constantinou M *see* Jakubowski L *et al*
Cooper M *see* Hart TC *et al*
Coovert DD *et al*. Does the survival motor neuron protein (SMN) interact with Bcl-2?: *letter*, **536**
Copin H *see* Ion A *et al*
Coppola B *see* Iolascon A *et al*
Corfield V *see* Moolman-Smook J *et al*
Cormier-Daire V *see* Faivre L *et al*
see Vuillaumier-Barrot S *et al*
Cormier-Daire V *et al*. Congenital disorders of glycosylation IIa cause growth retardation, mental retardation, and facial dysmorphism: *letter*, **875**

Genitopatellar syndrome: a new condition comprising absent patellae, scrotal hypoplasia, renal anomalies, facial dysmorphism, and mental retardation, **520**
Cornelio F *see* Barresi R *et al*
Correa-Cerro LS *et al*. The gene for branchio-oculo-facial syndrome does not colocalise to the *EYA1-4* genes: *letter*, **620**
Corris P *see* Thomson JR *et al*
Costa FF *see* Vasconcellos JPC *et al*
Costa M *et al*. Evaluation of the *HOX11L1* gene as a candidate for congenital disorders of intestinal innervation: *letter*, **e9**
Costa VP *see* Vasconcellos JPC *et al*
Costello JF *see* Frühwald MC *et al*
Coucourde G *see* Perfumo C *et al*
Cousin L *see* Manouvrier-Hanu S *et al*
Couturier A *see* Hutter P *et al*
Cox MC *see* Davies JF *et al*
Craen M *see* Mortier G *et al*
Craigen WJ *see* Kosaki K *et al*
Craufurd D *see* Harper PS *et al*
Crawford VLS *see* McIlroy SP *et al*
Cremers CWRJ *see* Pieke-Dahl S *et al*
Crolla JA *see* Cockwell AE *et al*
Crosby AH *see* Ion A *et al*
Cross SJ *et al*. The mutation spectrum in Holt-Oram syndrome: *letter*, **785**
Crozier A *see* Thomson JR *et al*
Cudry S *et al*. MPS II in females: molecular basis of two different cases: *letter*, **e29**
Cuer M *see* Vuillaumier-Barrot S *et al*
Cusano R *see* Costa M *et al*
Cusin V *see* Faivre L *et al*
Cusine D *see* Dean JCS *et al*
Cussenot O *see* Ion A *et al*
Cutting GR *see* Marcos I *et al*

Dabora SL *et al*. Characterisation of six large deletions in *TSC2* identified using long range PCR suggests diverse mechanisms including Alu mediated recombination: *letter*, **877**
Dagli *see* Kilinç *et al*
Dahia PM *et al*. Absence of germline mutations in *MINPP1*, a phosphatase encoding gene centromeric of *PTEN*, in patients with Cowden and Bannayan-Riley- Ruvalcaba syndrome without germline *PTEN* mutations: *letter*, **715**
Dahl N *see* Tentler D *et al*
Dake Y *see* Gao P-S *et al*
Dalakas MC *see* Park K-Y *et al*
Dalprà L *see* Villa N *et al*
Dalton A *see* Boyne J *et al*
Daniel S *see* Barresi R *et al*
Daruwala R *see* Stratakis CA *et al*
Das D *see* Reardon W *et al*
Da Silva R *see* de Lange RPJ *et al*
Dauwerse H *see* Eussen BHJ *et al*
Dauwerse HG *see* Petrij F *et al*
Davidson HR *see* Whiteford ML *et al*
Davidson R *see* Campbell H *et al*
Davies JF *et al*. 2157delG: a frequent mutation in *BRCA2* missed by PTT: *letter*, **e42**
Davis AC *see* Hutchin TP *et al*
see Parker MJ *et al*
Davison F *see* D'Cruz LG *et al*
Day INM *see* Heath KE *et al*
D'Cruz LG *et al*. Cytosine methylation confers instability on the cardiac troponin T gene in hypertrophic cardiomyopathy: *letter*, **e18**
Dean J *see* Campbell H *et al*
see Sargent CA *et al*
Dean JCS *see* Moore SJ *et al*
Dean JCS *et al*. Genetic registers in clinical practice: a survey of UK clinical geneticists: *letter*, **636**
Dearlove AM *see* Baralle D *et al*
Dearnaley DP *see* Forrest MS *et al*
de Bernabé D-V *see* Kilinç *et al*
de Braekeleer M *see* Rivard SR *et al*
de Córdoba SR *see* Kilinç *et al*
de Diego Otero Y *see* Willemsen R *et al*
Deeble J *see* Hutchin TP *et al*
Deeble VJ *et al*. The continuing failure to recognise Alström syndrome and further evidence of genetic homogeneity: *letter*, **219**
de la Chapelle A *see* Aktan-Collan K *et al*
see Desai DC *et al*
de Lange RPJ *et al*. Screening British CADASIL families for mutations in the *NOTCH3* gene: *letter*, **224**
de Lange W *see* Moolman-Smook J *et al*
Delatycki MB *et al*. Friedreich ataxia: an overview: *review*, **1**
Delhanty JDA *see* Hastings RJ *et al*
Dell'orto M *see* Villa N *et al*
de Lonlay P *see* Vuillaumier-Barrot S *et al*
Delozier-Blanchet C *see* Reardon W *et al*
de Marchi M *see* Bruttini M *et al*
de Martinville B *see* Manouvrier-Hanu S *et al*
Demir E *see* Dinçer P *et al*
see Özkinay F *et al*
Demirkol *see* Kilinç *et al*
Dennis N *see* Youings SA *et al*
Dennis NR *see* Marsh AJ *et al*
de Paepe A *see* Mortier G *et al*
see Mortier GR *et al*
see Nuytinck L *et al*
Depetris D *see* Cardoso C *et al*
Deprez PH *see* Chen J-M *et al*
de Ravel TJL *et al*. Idiopathic multicentric osteolysis presents early and is not linked to chromosome 18q21.1: *letter*, **e34**
de Rosa M *see* Kirsch S *et al*
Desai DC *et al*. Recurrent germline mutation in *MSH2* arises frequently de novo, **646**
de Salvo L *see* Montera M *et al*
Dessars B *see* Abramowicz MJ *et al*
de Villiers JNP *see* Thiart R *et al*
Devisme L *see* Manouvrier-Hanu S *et al*
de Vivo I *et al*. Novel germline mutations in the *PTEN* tumour suppressor gene found in women with multiple cancers, **336**
de Vries PJ, Bolton PF. Genotype-phenotype correlations in tuberous sclerosis: *letter*, **e3**
di Blasi C *see* Barresi R *et al*
Digilio MC *see* Tiberio G *et al*
Dijkstra PF *see* Terhal PA *et al*
di Lucca C *see* Eussen BHJ *et al*
Dinçer P *et al*. A cross section of autosomal recessive limb-girdle muscular dystrophies in 38 families, **361**
Ditchfield M *see* Halliday JL *et al*
Dixkens C *see* Knoblauch H *et al*
Doi K *see* Usami S-I *et al*
Dollinger MM *see* Graham AM *et al*
Domènech A *see* Nadal M *et al*
Donadini A *see* Montera M *et al*
Donald P *see* Fagan K *et al*
Donaldson B *see* Whiteford ML *et al*
Donnai D *see* Newman B *et al*
Dow D *see* Hes FJ *et al*
Dowe A *see* Forrest MS *et al*
Drira M *see* Masmoudi S *et al*
Dubourg O *see* Forissier J-F *et al*
Duggins A *see* Ellaway CJ *et al*
Dumez Y *see* Sanlaville D *et al*
Dupré T *see* Vuillaumier-Barrot S *et al*
Durand G *see* Cormier-Daire V *et al*
see Vuillaumier-Barrot S *et al*
Dutly F *see* Brecevic L *et al*
see Kotzot D *et al*
Dynan KB *see* McIlroy SP *et al*

Earl JW *see* Ellaway CJ *et al*
Easton D *see* Narain Y *et al*
Easton DF *see* Badzioch M *et al*
see Forrest MS *et al*
Eccles D *see* Hodgson SV *et al*
Eccles DM *et al*. Guidelines for a genetic risk based approach to advising women with a family history of breast cancer, **203**
Edery P *see* Morlé L *et al*
Edkins S *see* Mangion J *et al*
Edwards S *see* Badzioch M *et al*
Edwards SM *see* Forrest MS *et al*
Eeles R *see* Badzioch M *et al*
Eeles RA *see* Forrest MS *et al*
Egan J *see* Thomson JR *et al*
Eggermann K *see* Mergenthaler S *et al*
Eggermann T *see* Mergenthaler S *et al*
Egilsson V *see* Sigbjörnsdottir BI *et al*
Ekert H *see* Halliday JL *et al*
Eklöf O *see* Tentler D *et al*
Elgaied-Boulila A *see* Masmoudi S *et al*
Elinder G *see* Tentler D *et al*
Ellaway CJ *et al*. Cockayne syndrome associated with low CSF 5-hydroxyindole acetic acid levels: *letter*, **553**
Elles R *see* Davies JF *et al*
Elliott GC *see* Thomson JR *et al*
Elliott PM *see* D'Cruz LG *et al*
Ellis D *et al*. Low prevalence of germline *BRCA1* mutations in early onset breast cancer without a family history: *letter*, **792**
Elwyn G *et al*. Shared decision making and non-directiveness in genetic counselling, **135**
Emberger W *et al*. A de novo complex chromosomal rearrangement involving chromosomes 2, 3, and 10 associated with microcephaly and early onset spasticity: *letter*, **892**
Endo S *see* Obata K *et al*
Eng C *see* Borrego S *et al*
see Syngal S *et al*
see Zhou X-P *et al*
Eng C. Will the real Cowden syndrome please stand up: revised diagnostic criteria: *commentary*, **828**
Engel JR *et al*. Epigenotype-phenotype correlations in Beckwith-Wiedemann syndrome, **921**

Engel W *see* Blanck C *et al*
Engels S *see* Blanck C *et al*
Engels S *et al*. A *SALL1* mutation causes a branchio-oto-renal syndrome-like phenotype: *letter*, **458**
Eng C *see* Dahia PM *et al*
Ennis S *see* Youings SA *et al*
Enomoto T *see* Gao P-S *et al*
Esiri MM *see* Narain Y *et al*
Estivill X *see* Kilinç *et al*
see Nadal M *et al*
Eussen BHJ *et al*. An unbalanced submicroscopic translocation t(8;16)(q24.3;p13.3)pat associated with tuberous sclerosis complex, adult polycystic kidney disease, and hypomelanosis of Ito, **287**
Evans DGR *see* Baser ME *et al*
see Davies JF *et al*
see Desai DC *et al*
see Eccles DM *et al*
see Howell A *et al*
Evans DGR *et al*. Neurofibromatosis type 2: *review*, **897**
Use of MRI and audiological tests in presymptomatic diagnosis of type 2 neurofibromatosis (NF2): *letter*, **944**
Evans G *see* Ellis D *et al*
see Hodgson SV *et al*
Evans JG *see* Narain Y *et al*
Evans S *see* Boyne J *et al*
Eyre DR *see* Mortier GR *et al*

Faber J *et al*. Clinical variability of Stickler syndrome with a *COL2A1* haploinsufficiency mutation: implications for genetic counselling: *letter*, **318**
Fagan K *et al*. Fine molecular mapping of the 4p16.3 aneuploidy syndromes in four translocation families: *letter*, **449**
Faivre L *et al*. Exclusion of chromosome 9 helps to identify mild variants of acromesomelic dysplasia Maroteaux type, **52**
Farndon PA *see* Dean JCS *et al*
Fava M *see* Costa M *et al*
Fehaid EL *see* Meeks M *et al*
Felisari G *see* Barresi R *et al*
Felix R. Channelopathies: ion channel defects linked to heritable clinical disorders: *review*, **729**
Fenske C *see* Ion A *et al*
Ferec C *see* Chen J-M *et al*
see Chen J-M *et al*
Férec C *see* Rivard SR *et al*
Ferák V *see* Zatková A *et al*
Feráková E *see* Zatková A *et al*
Ferner R *see* John AM *et al*
Ferrans VJ *see* Park K-Y *et al*
Ferrer-Calvete J *see* Özkinay F *et al*
Ffrench-Constant C *see* Baralle D *et al*
Fiche M *see* Vaurs-Barrière C *et al*
Finkelstien S *see* Lev D *et al*
Firatli E *see* Hart PS *et al*
see Hart TC *et al*
Fitzpatrick DR *see* Dean JCS *et al*
Flanagan AM *see* Mangion J *et al*
Flint J *see* Knight SJL and Flint J
Flinter F *see* Warburg M *et al*
Flinter FA *see* Plant KE *et al*
Flynn DM *see* Sidwell RU *et al*
Fois A *see* Eussen BHJ *et al*
Fok T-F *see* Lam C-W *et al*
Fokstuen S, Schinzel A. Unilateral lobar pulmonary agenesis in sibs: *letter*, **557**
Fontes M *see* Cardoso C *et al*
Forabosco P *see* Costa M *et al*
Forissier J-F *et al*. First description of germline mosaicism in familial hypertrophic cardiomyopathy, **132**
Forrest MS *et al*. No evidence of germline *PTEN* mutations in familial prostate cancer: *letter*, **210**
Forrest SM *see* Delatycki MB *et al*
Forrest SM *see* Massie J *et al*
Forshaw K *see* Massie J *et al*
Fortnum HM *see* Parker MJ *et al*
Foulkes WD *see* Badzioch M *et al*
see Metcalfe KA *et al*
Fox EA *see* Syngal S *et al*
France L *see* Gray J *et al*
Francis PJ *et al*. The genetics of childhood cataract: *review*, **481**
Franz D *see* Dabora SL *et al*
Freiman J *see* Koorey D *et al*
Frets PG *see* Lodder LN *et al*
Frey H *see* Mattila KM *et al*
Friedl W *see* Gohlke BC *et al*
Friedman JM *see* Szudek J *et al*
Friedrich U *see* Bugge M *et al*
Froissart R *see* Cudry S *et al*
Fruhwald MC *et al*. Gene amplification in PNETs/medulloblastomas: mapping of a novel amplified gene within the *MYCN* amplicon, **501**
Fryns JP *see* de Ravel TJL *et al*
Fryns J-P, Aftimos S. New MR/MCA syndrome with distinct facial appearance and general habitus, broad and webbed neck, hypoplastic inverted nipples, epilepsy, and pachygyria of the frontal lobes: *letter*, **460**
Fukai K *see* Saitoh S *et al*
Fukami M *see* Gohlke BC *et al*
Fukuda T *see* Obata K *et al*
Fukuzawa R *see* Kikuchi H *et al*
Fung VS *see* Ellaway CJ *et al*
Furlong RA *see* Keramatipour M *et al*
see Narain Y *et al*

Gaffney D *see* Thiart R *et al*
Gallet S *see* Bahuau M *et al*
Gao P-S *et al*. Variants of *STAT6* (signal transducer and activator of transcription 6) in atopic asthma: *letter*, **380**
Gao X *see* Borrego S *et al*
see Desai DC *et al*
see Zhou X-P *et al*
Garber JE *see* Syngal S *et al*
Gardiner RM *see* Meeks M *et al*
Gardiner C-A *see* Cross SJ *et al*
Gardner RJM *see* Savarirayan R *et al*
see Voullaire L *et al*
Gaunt L *see* Slavotinek AM *et al*
Gécz J. *FMR3* is a novel gene associated with *FRAXE* CpG island and transcriptionally silent in *FRAXE* full mutations, **782**
Georgantopoulou A *see* Berg JN *et al*
Gerrard M *see* Pritchard-Jones K *et al*
Gershoni-Baruch R *see* Janssens K *et al*
Gertig DM *see* de Vivo I *et al*
Giannotti A *see* Tiberio G *et al*
Gibbons B *see* Cockwell AE *et al*
see Sidwell RU *et al*
Gibbs JSR *see* Thomson JR *et al*
Gilbert B *see* Faivre L *et al*
Giles RH *see* Petrij F *et al*
Giles G *see* Badzioch M *et al*
Gillespie J *see* Evans DGR *et al*
Gillessen-Kaesbach G *see* Wieczorek D *et al*
Gimm O *see* Borrego S *et al*
see Dahia PM *et al*
see Zhou X-P *et al*
Ginns EI *see* Stone DL *et al*
Glavač D *see* Ravnik-Glavač M *et al*
Glaze D *see* Potocki L *et al*
Glover G *see* Özkinay F *et al*
Glover S *see* Moore SJ *et al*
Gnoinski W *see* Faber J *et al*
Godet J *see* Morlé L *et al*
Godfrey M *see* Robinson PN and Godfrey M
Goebel HH *see* Park K-Y *et al*
Goh DLM *et al*. Dysmorphic sibs trisomic for the region 6q22.1→6q23.3: *letter*, **889**
Gohlke BC *et al*. Interstitial deletion in Xp22.3 is associated with X linked ichthyosis, mental retardation, and epilepsy, **600**
Goldfarb LG *see* Park K-Y *et al*
Goldgar D *see* Badzioch M *et al*
Goodman BK *see* Praphanphoj V *et al*
Goossens M *see* Abramowicz MJ *et al*
Gordon L *see* Tentler D *et al*
Gordon RD *see* Lafferty AR *et al*
Gorin MB *see* Pieke-Dahl S *et al*
Gorlin RJ. Of palms, soles, and gums: *commentary*, **81**
Goss AN *see* Mangion J *et al*
Goudie D *see* Campbell H *et al*
Graham AM *et al*. Identification of novel alleles at a polymorphic microsatellite repeat region in the human *NRAMP1* gene promoter: analysis of allele frequencies in primary biliary cirrhosis: *letter*, **150**
Grandchamp B *see* Vuillaumier-Barrot S *et al*
Grant ME *see* Gregory CA *et al*
Gray J *see* Brain K *et al*
see Elwyn G *et al*
Gray J *et al*. A model protocol evaluating the introduction of genetic assessment for women with a family history of breast cancer, **192**
Graziani M *see* Tiberio G *et al*
Graziano C *see* Kilinç *et al*
Green EK *et al*. Detailed mapping of a congenital heart disease gene in chromosome 3p25, **581**
Green PM *see* Plant KE *et al*
Greenman J *see* Ellis D *et al*
Greer WL *see* Beresford RG *et al*
Gregory CA *et al*. Equal expression of type X collagen mRNA from mutant and wild type *COL10A1* alleles in growth plate cartilage from a patient with metaphyseal chondrodysplasia type Schmid: *letter*, **627**
Grioni D *see* Villa N *et al*
Grossmann JG *see* Howell A *et al*
Guañabens N *see* Janssens K *et al*
Guanti G *see* Montera M *et al*

Guggenheim JA *et al.* The heritability of high myopia: a reanalysis of Goldschmidt's data: *letter*, **227**
Guldberg P *see* Zschocke J *et al*
Gustavsson P *see* Tentler D *et al*
Gutkowska A *see* Kotzot D *et al*

Hachicha S *see* Masmoudi S *et al*
Hainque B *see* Forissier J-F *et al*
Haites NE *see* Hodgson SV *et al*
Hall K *see* Lynch SA *et al*
Halley DJJ *see* Eussen BHJ *et al*
Halliday JL *et al.* Hemiplegic cerebral palsy and the factor V Leiden mutation: *letter*, **787**
Haltia T *see* Nopola-Hemmi J *et al*
Hamam HD *see* Meeks M *et al*
Hamasaki N *see* Gao P-S *et al*
Hampel H *see* Westman J *et al*
Hampson K *et al.* Mutations in the *MECP2* gene in a cohort of girls with Rett syndrome: *letter*, **610**
Hankinson SE *see* de Vivo I *et al*
Hansen J *see* Bugge M *et al*
Hansford JR, Mulligan LM. Multiple endocrine neoplasia type 2 and *RET*: from neoplasia to neurogenesis: *review*, **817**
Hardouin A *see* Vaurs-Barrière C *et al*
Harper A *see* Engel JR *et al*
Harper P *see* Brain K *et al*
see Gray J *et al*
Harper PS *et al.* Ten years of presymptomatic testing for Huntington's disease: the experience of the UK Huntington's Disease Prediction Consortium, **567**
Harper P *see* McEntagart M *et al*
Harrington P *see* Seltmann M *et al*
Harrison DJ *see* Graham AM *et al*
Hart PS *see* Hart TC *et al*
see Hart TC *et al*
Hart PS *et al.* Identification of cathepsin C mutations in ethnically diverse Papillon-Lefèvre syndrome patients, **927**
Hart TC *see* Hart PS *et al*
Hart TC *et al.* Haim-Munk syndrome and Papillon-Lefèvre syndrome are allelic mutations in cathepsin C, **88**
Localisation of a gene for prepubertal periodontitis to chromosome 11q14 and identification of a cathepsin C gene mutation, **95**
Hasegawa T *see* Lichtner P *et al*
Hasnain SS *see* Howell A *et al*
Hastings RJ *et al.* Deletion and duplication of the adenomatous polyposis coli gene resulting from an interchromosomal insertion involving 5(q22q23.3) in the father: *letter*, **141**
Hastings RJ *see* Sidwell RU *et al*
Hata J-I *see* Kikuchi H *et al*
Hatchwell E *see* Browne CE *et al*
Haug K *see* Gohlke BC *et al*
Hauschild M *see* Scharfe C *et al*
Haverkamp F *see* Gohlke BC *et al*
Hazan J *see* Lindsey JC *et al*
Heath KE *et al.* I705 variant in the low density lipoprotein receptor gene has no effect on plasma cholesterol levels: *letter*, **713**
Universal primer quantitative fluorescent multiplex (UPQFM) PCR: a method to detect major and minor rearrangements of the low density lipoprotein receptor gene, **272**
Heathcote K *et al.* A connexin 26 mutation causes a syndrome of sensorineural hearing loss and palmoplantar hyperkeratosis (MIM 148350), **50**
Heidemann PH *see* Scharfe C *et al*
Heimdal K *see* Badzioch M *et al*
Heins YM *see* Nieuwint AWM *et al*
Heller A *et al.* Molecular cytogenetic characterisation of partial trisomy 9q in a case with pyloric stenosis and a review, **529**
Helszer Z *see* Jakubowski L *et al*
Henderson S *et al.* Germline and somatic mosaicism in achondroplasia: *letter*, **956**
Hennekam RCM *see* Kelley RI and Hennekam RCM
see Petrij F *et al*
see Waterham HR *et al*
Henske EP *see* Astrinidis A *et al*
Héon E *see* Martin SN *et al*
Heppell-Parton AC *see* Seltmann M *et al*
Hernández A *see* Borrego S *et al*
Hes FJ *et al.* Cryptic von Hippel-Lindau disease: germline mutations in patients with haemangioblastoma only, **939**
Hetet G *see* Vuillaumier-Barrot S *et al*
Higenbottam T *see* Thomson JR *et al*
Higgins MJ *see* Engel JR *et al*
Hillebrand G *et al.* DiGeorge syndrome with discordant phenotype in monozygotic twins: *letter*, **e23**
Hindmarsh P *see* Reardon W *et al*
Hitchins M *see* Wakeling EL *et al*
Hitchins MP *see* Wakeling EL *et al*
Hodgson S *see* Ellis D *et al*
Hodgson SV *et al.* A survey of the current clinical facilities for the management of familial cancer in Europe, **605**
Hodson SA *see* Guggenheim JA *et al*
Hoffmann GF *see* Zschocke J *et al*
Hoffs MS *see* Thiart R *et al*
Hole R *see* Turnpenny PD and Hole R
Holloway S *see* Campbell H *et al*
Holvoet M *see* de Ravel TJL *et al*
Homfray T *see* Wuyts W *et al*
Honda M *see* Kikuchi H *et al*
Honour JW *see* Reardon W *et al*
Hoodfar E *see* Metcalfe KA *et al*
Hoogendijk CF *see* Thiart R *et al*
Hooper ML *see* Young IE *et al*
Hoover D *see* Smith SD *et al*
Hopkin JM *see* Gao P-S *et al*
Hopper JL *see* Badzioch M *et al*
Horiuchi I *see* Obata K *et al*
Hoshino T *see* Usami S-I *et al*
Houlston RS *see* Lucassen AM and Houlston RS
Houlston RS, Tomlinson IPM. Detecting low penetrance genes in cancer: the way ahead: *review*, **161**
Howell A *et al.* Can hair be used to screen for breast cancer?: *letter*, **297**
Howie SEM *see* Graham AM *et al*
Hsu E *see* Wang J *et al*
Hubler A *see* Heller A *et al*
Huggard P *see* Lafferty AR *et al*
Hui J *see* Lam C-W *et al*
Huiping C *see* Sigbjörnsdottir BI *et al*
Humbert M *see* Thomson JR *et al*
Hummel M *see* Ndah BV *et al*
Humphreys L *et al.* Why patients do not attend for their appointments at a genetics clinic: *letter*, **810**
Humphries SE *see* Heath KE *et al*
see Heath KE *et al*
Huner *see* Kilinç *et al*
Hunter AGW *see* Armour CM *et al*
see Humphreys L *et al*
Hunter DJ *see* de Vivo I *et al*
Hurko O *see* Wright MJ *et al*
Hurles ME *see* Blanco P *et al*
Hurst JA *see* Lynch SA *et al*
Hutchin TP *et al.* A novel mutation in the mitochondrial $tRNA^{Ser(UCN)}$ gene in a family with non-syndromic sensorineural hearing impairment, **692**
Hutter P *et al.* Two common forms of the human *MLH1* gene may be associated with functional differences, **776**

Imaizumi K *see* Mirza G *et al*
Imhof SM *see* Scheffer H *et al*
Ingvarsson S *see* Sigbjörnsdottir BI *et al*
Iolascon A *et al.* Crigler-Najjar syndrome type II resulting from three different mutations in the bilirubin uridine 5'-diphosphate-glucuronosyltransferase (*UGT1A1*) gene: *letter*, **712**
Ion A *et al.* Abnormal sex differentiation and multiple congenital abnormalities in a subject harbouring an apparently balanced (6;8) translocation: *letter*, **972**
Detailed mapping, mutation analysis, and intragenic polymorphism identification in candidate Noonan syndrome genes *MYL2*, *DCN*, *EPS8*, and *RPL6* : *letter*, **884**
Ishii M *see* Usami S-I *et al*
Ito J *see* Usami S-I *et al*
Ito S *see* Kikuchi H *et al*
Iwanaga R *see* Obata K *et al*
Iwasaki S *see* Usami S-I *et al*
Izatt L *see* Ellis D *et al*
Izuhara K *see* Gao P-S *et al*

Jackson A *see* Deeble VJ *et al*
see Slavotinek AM *et al*
Jackson C *see* McEntagart M *et al*
Jacobi C *see* Plaschke J *et al*
Jacobs C *see* Ellis D *et al*
Jacobs P *see* Youings SA *et al*
Jaksch M *see* Scharfe C *et al*
Jakubów-Durska K *see* Stankiewicz P *et al*
Jakubowski L *et al.* Xp;Yp translocation inherited from the father in an *SRY*, *RBM*, and *TSPY* positive true hermaphrodite with oligozoospermia: *letter*, **e28**
Janssen AJM *see* Scharfe C *et al*
Janssens K *et al.* Localisation of the gene causing diaphyseal dysplasia Camurati-Engelmann to chromosome 19q13, **245**
Järvinen H *see* Roth S *et al*
Jeanpierre C *see* Manouvrier-Hanu S *et al*
Jeffery S *see* Bell R *et al*
Jeffery S *see* Ion A *et al*
Jenkins JPR *see* Evans DGR *et al*
Jensen G *see* Bugge M *et al*
Jensen PKA *see* Bugge M *et al*
Jeziorowska A *see* Jakubowski L *et al*
Jobling MA *see* Blanco P *et al*
Johanning GL *et al.* Comorbidity of 5,10-methylenetetrahydrofolate reductase and methionine synthase gene polymorphisms and risk for neural tube defects: *letter*, **949**

Johansson M *see* Roth S *et al*
John AM *et al.* A search for evidence of somatic mutations in the *NF1* gene, **44**
Johnson CA. Chromatin modification and disease: *review*, **905**
Johnson EI *et al.* 5p14 deletion associated with microcephaly and seizures, **125**
Johnston KE *see* Johanning GL *et al*
Jones EA *et al. JAGGED1* expression in human embryos: correlation with the Alagille syndrome phenotype, **658**
Jonveaux P *see* Cross SJ *et al*
Joyce JA *see* Catchpoole D *et al*
Jozwiak S *see* Dabora SL *et al*
Just W *see* Correa-Cerro LS *et al*

Kaasik K *see* Teder M *et al*
Kacet N *see* Manouvrier-Hanu S *et al*
Kádasi L *see* Zatková A *et al*
Kääriäinen H *see* Aktan-Collan K *et al*
Kale G *see* Dinçer P *et al*
Kalimo H *see* Mattila KM *et al*
Kamath R *see* Ellaway CJ *et al*
Kanbi L *see* Howell A *et al*
Karaduman A *see* Dinçer P *et al*
Karakoç *see* Kilinç *et al*
Karaüzüm SB *see* Kotzot D *et al*
Karbani G *see* Deeble VJ *et al*
Kashork CD *see* Potocki L *et al*
Kassab A *see* Masmoudi S *et al*
Kassab I *see* Masmoudi S *et al*
Katzin N *see* Lev D *et al*
Kaułzewski B *see* Jakubowski L *et al*
Kawai M *see* Gao P-S *et al*
Kayserili H *see* Nuytinck L *et al*
Kelley PM *see* Abe S *et al*
see Pieke-Dahl S *et al*
see Smith SD *et al*
Kelley RI, Hennekam RCM. The Smith-Lemli-Opitz syndrome: *review*, **321**
Kelly J *see* Forrest MS *et al*
Kelly T *see* Reardon W *et al*
Kelly TE *et al.* Confirmation of the assignment of the Sanjad-Sakati (congenital hypoparathyroidism) syndrome (OMIM 241410) locus to chromosome lq42–43: *letter*, **63**
Kenmochi N *see* Ion A *et al*
Kennerknecht I *see* Correa-Cerro LS *et al*
Kenyon JB *see* Smith SD *et al*
Keramatipour M *et al.* The *ACE* I allele is associated with increased risk for ruptured intracranial aneurysms, **498**
Kere J *see* Nopola-Hemmi J *et al*
Khare L *see* Astrinidis A *et al*
Kidd A *see* Sargent CA *et al*
Kiechle M *see* Weimer J *et al*
Kikuchi H *et al.* Constitutional *WT1* mutations correlate with clinical features in children with progressive nephropathy: *letter*, **698**
Kilimann MW *see* Burwinkel B *et al*
Kilinç *et al.* Alkaptonuria in Italy: polymorphic haplotype background, mutational profile, and description of four novel mutations in the homogentisate 1,2-dioxygenase gene: *letter*, **309**
Kimberling WJ *see* Abe S *et al*
see Pieke-Dahl S *et al*
Kimura A *see* Obata K *et al*
King LM *see* Mortier GR *et al*
Kingston H *see* Slavotinek A and Kingston H
Kingston H *see* Dean JCS *et al*
King-Underwood L *see* Pritchard-Jones K *et al*
Kinmond S *see* Whiteford ML *et al*
Kirkpatrick P *see* Keramatipour M *et al*
Kirov G *see* Guggenheim JA *et al*
Kirsch S *et al.* FISH deletion mapping defines a single location for the Y chromosome stature gene, *GCY*, **593**
Klaassen T *see* Teder M *et al*
Klijn JGM *see* Lodder LN *et al*
Klopstock T *see* Scharfe C *et al*
Klose R *see* Martin SN *et al*
Knight SJL, Flint J. Perfect endings: a review of subtelomeric probes and their use in clinical diagnosis: *review*, **401**
Knoblauch H *et al.* Clinical and molecular cytogenetic studies of a large de novo interstitial deletion 16q11.2-16q21 including the putative transcription factor gene *SALL1* : *letter*, **389**
Knowles MA *see* Forrest MS *et al*
Kohlhase J *see* Blanck C *et al*
see Engels S *et al*
see Knoblauch H *et al*
Koivula T *see* Mattila KM *et al*
Kolodner RD *see* Syngal S *et al*
Komajda M *see* Forissier J-F *et al*
Komiyama S *see* Usami S-I *et al*
Komune S *see* Usami S-I *et al*
Kondo I *see* Obata K *et al*
König R *see* Lichtner P *et al*
Koorey D *et al.* Appendiceal carcinoma complicating adenomatous polyposis in a young woman with a de novo constitutional reciprocal translocation t(5;8)(q22;p23.1): *letter*, **71**
Ko C-H *see* Lam C-W *et al*
Korenberg JR *see* Slavotinek AM *et al*
Korneluk Y *see* Humphreys L *et al*
Körner H *see* Knoblauch H *et al*
Korngut LM *see* Onochie CI *et al*
Kosaki K *et al.* Diagnosis of maternal uniparental disomy of chromosome 7 with a methylation specific PCR assay: *letter*, **e19**
Kosaki R *see* Kosaki K *et al*
Kotze MJ *see* Thiart R *et al*
Kotzot D *et al.* Parental origin and mechanisms of formation of cytogenetically recognisable de novo direct and inverted duplications, **281**
Krahe R *see* Frühwald MC *et al*
Krajewska-Walasek M *see* Kotzot D *et al*
Kralewski M *see* Coovert DD *et al*
Krause M *see* Wieczorek D *et al*
Kremenski I *see* Özkinay F *et al*
Kremer H *see* Ion A *et al*
Kristoffersson U *see* Bugge M *et al*
Kroes HY *see* Blanck C *et al*
Kroisel PM *see* Emberger W *et al*
Kroisel PM *et al.* Skin pigmentary anomalies in a mosaic form of partial tetrasomy 3q: *letter*, **723**
Kubo T *see* Usami S-I *et al*
Kucherlapati RS *see* Ion A *et al*
Kuijpers GMC *see* Terhal PA *et al*
Kumar D *see* Reardon W *et al*
Kunkler IH *see* Young IE *et al*
Kurian KM *see* Young IE *et al*
Kuwajima K *see* Obata K *et al*
Kwiatkowski DJ *see* Dabora SL *et al*
Kwong NS *see* Lam FWF *et al*
Kytölä S *see* Poyhonen M *et al*

Laccone F *see* Maiwald R *et al*
Lachman RS *see* Mortier GR *et al*
Lacombe D *see* Bahuau M *et al*
Lafferty AR *et al.* A novel genetic locus for low renin hypertension: familial hyperaldosteronism type II maps to chromosome 7 (7p22), **831**
Laippala P *see* Mattila KM *et al*
Lalloo F *see* Ellis D *et al*
Lalloo FI *see* Davies JF *et al*
Lam C-W *et al.* Spectrum of mutations in the *MECP2* gene in patients with infantile autism and Rett syndrome: *letter*, **e41**
Lam FWF *et al.* Proximal 10q trisomy: a new case with anal atresia: *letter*, **e24**
Lam STS *see* Lam FWF *et al*
Lam W *see* Catchpoole D *et al*
Lamireau T *see* Bahuau M *et al*
Laplace-Marieze V *see* Vaurs-Barrière C *et al*
Latif F *see* Green EK *et al*
see Hampson K *et al*
Lato E *see* Stankiewicz P *et al*
Latour P *see* Morlé L *et al*
Laurendeau I *see* Bahuau M *et al*
Lawson JT *see* McIlroy SP *et al*
Le TT *see* Coovert DD *et al*
le Bail B *see* Bahuau M *et al*
Leblanc J-P *see* Rivard SR *et al*
Leblanc G *see* Badzioch M *et al*
Lederer G *see* Schmidt H *et al*
Ledeuil C *see* Forissier J-F *et al*
Lees J *see* Portwine C *et al*
Leggatt V *et al.* The psychological impact of a cancer family history questionnaire completed in general practice: *letter*, **470**
Legius E *see* de Ravel TJL *et al*
see López-Correa C *et al*
Lehesjoki A-E *see* Nopola-Hemmi J *et al*
Leheup B *see* Cross SJ *et al*
Lehtimäki T *see* Mattila KM *et al*
Leisti J *see* Poyhonen M *et al*
Le Merrer M *see* Cormier-Daire V *et al*
see Cormier-Daire V *et al*
see Faivre L *et al*
Lench N *see* Deeble VJ *et al*
Lench NJ *see* Hutchin TP *et al*
Lequien P *see* Manouvrier-Hanu S *et al*
Lerman-Sagie T *see* Lev D *et al*
see Lev D *et al*
Lerone M *see* Costa M *et al*
Leroy JG *see* Mortier G *et al*
Lesourd A *see* Ion A *et al*
Lev D *et al.* CHARGE association in a child with de novo chromosomal aberration 46,X,der(X)t(X;2)(p22.1;q33) detected by spectral karyotyping: *letter*, **e47**
Progressive neurological deterioration in a child with distal arthrogryposis and whistling face: *letter*, **231**
Levin AV *see* Martin SN *et al*

Levine M *see* Stratakis CA *et al*
Levy M, Richard S. Attitudes of von Hippel-Lindau disease patients towards presymptomatic genetic diagnosis in children and prenatal diagnosis: *letter*, **476**
Lewis C *see* Youings SA *et al*
Li FP *see* Portwine C *et al*
Lichtner P *et al.* An HDR (hypoparathyroidism, deafness, renal dysplasia) syndrome locus maps distal to the DiGeorge syndrome region on 10p13/14, **33**
Liede A *see* Metcalfe KA *et al*
Liede A *et al.* Delineation of a new syndrome: clustering of pyloric stenosis, endometriosis, and breast cancer in two families: *letter*, **794**
Liehr T *see* Heller A *et al*
Lim C *see* Harper PS *et al*
Lin JP *see* Lafferty AR *et al*
Lina-Granade G *see* Morlé L *et al*
Lindsay S *see* Lynch SA *et al*
Lindsey JC *et al.* Mutation analysis of the spastin gene (*SPG4*) in patients with hereditary spastic paraparesis, **759**
Lips CJM *see* Hes FJ *et al*
Li QY *see* Cross SJ *et al*
Litvak S *see* Park K-Y *et al*
Lloyd DJ *see* Moore SJ *et al*
Lo IFM *see* Lam C-W *et al*
Lo YMD *see* Lam C-W *et al*
Lockman JC *see* Desai DC *et al*
Lodder LN *et al.* Attitudes towards termination of pregnancy in subjects who underwent presymptomatic testing for the *BRCA1*/*BRCA2* gene mutation in The Netherlands: *letter*, **883**
Löfgren A *see* van Goethem G *et al*
Lombardi G *see* Kirsch S *et al*
López-Alonso M *see* Borrego S *et al*
López-Correa C *et al. NF2* gene deletion in a family with a mild phenotype: *letter*, 75
Lotfazar M *see* Hart PS *et al*
Loughlin J *see* Henderson S *et al*
Loukola A *see* Roth S *et al*
see Wheeler JMD *et al*
Loyd JE *see* Thomson JR *et al*
Lu X *see* Hart PS *et al*
Lubetzki J *see* Ion A *et al*
Lubs HA *see* Cabezas DA *et al*
Lucassen AM, Houlston RS. Clinical geneticists' attitudes and practice towards testing for breast cancer susceptibility genes: *letter*, **157**
Ludman MD *see* Beresford RG *et al*
Luleci G *see* Kotzot D *et al*
Lundsteen C *see* Bugge M *et al*
Lunt PW *see* Stalker DJ *et al*
Lupski JR *see* Potocki L *et al*
Lusher ME *see* Lindsey JC *et al*
Lutz Y *see* Cardoso C *et al*
Lynch SA *et al.* Autosomal dominant sacral agenesis: Currarino syndrome: *review*, **561**
Two further cases of Sener syndrome: frontonasal dysplasia and dilated Virchow-Robin spaces: *letter*, **466**
Lyonnet S *see* Bahuau M *et al*
see Cormier-Daire V *et al*
see Sanlaville D *et al*
Lyonnet S *see* Cross SJ *et al*

Maaswinkel-Mooy PD *see* Petrij F *et al*
McCall S *see* Ellis D *et al*
McConnell RS *see* Keramatipour M *et al*
Macdermot K *see* Aldred MA *et al*
see Ellis D *et al*
McDermott CJ *see* Lindsey JC *et al*
MacDonald F *see* Hes FJ *et al*
McDowell M *see* Savarirayan R *et al*
McEntagart M *et al.* Distal spinal muscular atrophy with vocal cord paralysis (dSMA-VII) is not linked to the *MPD2* locus on chromosome 5q31: *letter*, **e14**
McGaughran J *see* Engels S *et al*
McGleenon BM *see* McIlroy SP *et al*
Machado RD *see* Thomson JR *et al*
McIlroy SP *et al.* Butyrylcholinesterase K variant is genetically associated with late onset Alzheimer's disease in Northern Ireland, **182**
McIntosh I *see* Wright MJ *et al*
Mackay J *see* Eccles DM *et al*
see Leggatt V *et al*
McKechnie N *see* Youings SA *et al*
McKee S *see* Hes FJ *et al*
see Reardon W *et al*
McKeehan WL *see* Reardon W *et al*
McKenna WJ *see* D'Cruz LG *et al*
Mackenzie MAF *see* Young IE *et al*
Mackinnon H *see* Hastings RJ *et al*
Macleod R *see* Evans DGR *et al*
McMahon R *see* Hes FJ *et al*
Madan K *see* Nieuwint AWM *et al*
Maher ER *see* Catchpoole D *et al*
see Desai DC *et al*
see Engel JR *et al*
see Green EK *et al*
see Hes FJ *et al*
see Webster AR *et al*
see Woodward EK *et al*
Mainardi PC *see* Perfumo C *et al*
Maire I *see* Cudry S *et al*
Maiwald R *et al.* No evidence for imprinting in distal 18q: *letter*, **152**
Majewski F *see* Wieczorek D *et al*
Malcolm S *see* Reardon W *et al*
Malerba G *see* Patuzzo C *et al*
Maliszewska C *see* Green EK *et al*
Malkin D *see* Portwine C *et al*
Mancini A *see* Montera M *et al*
Mandel A *see* Tentler D *et al*
Mangion J *et al.* Familial craniofacial fibrous dysplasia: absence of linkage to *GNAS1* and the gene for cherubism: *letter*, **e37**
Mannoni A *see* Kilinç *et al*
Manouvrier-Hanu S *et al.* Sex reversal and diaphragmatic hernia in phenotypically female sibs with normal XY chromosomes: *letter*, 315
Man NT *see* Coovert DD *et al*
Mansel R *see* Brain K *et al*
see Gray J *et al*
Mansour S *see* Bell R *et al*
Mao X-Q *see* Gao P-S *et al*
Marazita ML *see* Hart TC *et al*
Marchese C *see* Montera M *et al*
Marcos I *et al.* Mutation analysis of *GABRR1* and *GABRR2* in autosomal recessive retinitis pigmentosa (*RP25*): *letter*, **e5**
Mareni C *see* Montera M *et al*
Mariman E *see* Ion A *et al*
Marinescu RC *see* Johnson EI *et al*
Markham AF *see* Hutchin TP *et al*
Marks JJ *see* Hart PS *et al*
Maroteaux P *see* Faivre L *et al*
Marsh AJ *et al.* Interstitial deletion of chromosome 17 (del(17)(q22q23.3 confirms a link with oesophageal atresia: *letter*, **701**
Marshall R *see* Hart PS *et al*
Marsh DJ *see* Dahia PM *et al*
Marteau TM *see* Leggatt V *et al*
Martin DM *et al.* Characterisation and genetic mapping of a new X linked deafness syndrome, **836**
Martin SN *et al.* Molecular characterisation of congenital glaucoma in a consanguineous Canadian community: a step towards preventing glaucoma related blindness, **422**
Martinati LC *see* Patuzzo C *et al*
Martinez M-J *see* Kotzot D *et al*
Martin J-J *see* van Goethem G *et al*
Martucciello G *see* Costa M *et al*
Masmoudi S *et al.* Determination of the frequency of connexin26 mutations in inherited sensorineural deafness and carrier rates in the Tunisian population using DGGE: *letter*, **e39**
Massie J *et al.* The relationship between neonatal immunoreactive trypsinogen, ΔF508, and IVS8-5T: *letter*, **629**
Mateu E *see* Sánchez F *et al*
Mathew CG *see* Ellis D *et al*
Matiaszuk N *see* Watkins D *et al*
Matsuishi T *see* Obata K *et al*
Matsumoto M *see* Yamamoto T *et al*
Matsuo N *see* Kosaki K *et al*
Mattei M-G *see* Cardoso C *et al*
Matthijs G *see* de Ravel TJL *et al*
Mattila KM *et al.* Dipeptidyl carboxypeptidase 1 (*DCP1*) and butyrylcholinesterase (*BCHE*) gene interactions with the apolipoprotein E ε4 allele as risk factors in Alzheimer's disease and in Parkinson's disease with coexisting Alzheimer pathology, **766**
Matušek J *see* Zatková A *et al*
Maugard CM *see* Vaurs-Barrière C *et al*
Mazurczak T *see* Stankiewicz P *et al*
Mazzucco G *see* Bruttini M *et al*
Mecklin J-P *see* Aktan-Collan K *et al*
see Roth S *et al*
Meeks M *et al.* A locus for primary ciliary dyskinesia maps to chromosome 19q, **241**
Megarbane A *see* Faivre L *et al*
Mehdi SQ *see* Bessant DAR *et al*
Meijers-Heijboer EJ *see* Lodder LN *et al*
Meinecke P *see* Wieczorek D *et al*
Meiser B *et al.* Attitudes to genetic testing for breast cancer susceptibility in women at increased risk of developing hereditary breast cancer: *letter*, **472**
Meitinger T *see* Lichtner P *et al*
see Scharfe C *et al*
Melo MB *see* Vasconcellos JPC *et al*
Meloni A *see* Iolascon A *et al*
Meloni I *see* Bruttini M *et al*

Mercier B *see* Chen J-M *et al*
Mergenthaler S *et al*. Exclusion of a disease relevant role of *PAX4* in the aetiology of Silver-Russell syndrome: screening for mutations and determination of imprinting status: *letter*, **e44**
Metcalfe KA *et al*. An evaluation of needs of female *BRCA1* and *BRCA2* carriers undergoing genetic counselling, **866**
Metspalu A *see* Teder M *et al*
Michalec MD *see* Hart PS *et al*
see Hart TC *et al*
see Hart TC *et al*
Michaud D *see* Onochie CI *et al*
Micutková L *see* Zatková A *et al*
Mignon C *see* Cardoso C *et al*
Migone N *see* Janssens K *et al*
Mikhail G *see* Thomson JR *et al*
Mikiewicz-Sygula D *see* Jakubowski L *et al*
Millat G *see* Yamamoto T *et al*
Miller GJ *see* Heath KE *et al*
Milot M *see* Rivard SR *et al*
Miny P *see* Kotzot D *et al*
Mirza G *et al*. Partial trisomy 22 in a liveborn resulting from a rearrangement between chromosomes 6 and 22: *letter*, **e22**
Miserez AR *see* Zumsteg U *et al*
Mitchell M *see* Liede A *et al*
Mitchison HM *see* Meeks M *et al*
Miyaki M *see* Desai DC *et al*
Modesto C *see* Patiño-Garcia A *et al*
Mohammed S *see* Ellis D *et al*
Moll AC *see* Scheffer H *et al*
Möller CG *see* Pieke-Dahl S *et al*
Moller P *see* Badzioch M *et al*
see Hodgson SV *et al*
Monk D *see* Wakeling EL *et al*
Montera M *et al*. Mutational germline analysis of *hMSH2* and *hMLH1* genes in early onset colorectal cancer patients: *letter*, **e7**
Montgomery T *see* Moore SJ *et al*
Montgomery K *see* Ion A *et al*
Moolman-Smook J *et al*. Expression of HCM causing mutations: lessons learnt from genotype-phenotype studies of the South African founder *MYH7* A797T mutation: *letter*, **951**
Moore AT *see* Francis PJ *et al*
see Yates JRW and Moore AT
Moore GE *see* Wakeling EL *et al*
see Wakeling EL *et al*
Moore S *see* Sargent CA *et al*
Moore SJ *et al*. A clinical study of 57 children with fetal anticonvulsant syndromes, **489**
Moore AT *see* Webster AR *et al*
Moore IE *see* Cockwell AE *et al*
Mora M *see* Barresi R *et al*
Moraine C *see* Forissier J-F *et al*
Morandi L *see* Barresi R *et al*
Morgan NV *see* Thomson JR *et al*
Mori K *see* Obata K *et al*
Morichon-Delvallez N *see* Sanlaville D *et al*
Morlé L *et al*. A novel C202F mutation in the connexin26 gene (*GJB2*) associated with autosomal dominant isolated hearing loss, **368**
Morrison P *see* Hodgson SV *et al*
Morris GE *see* Coovert DD *et al*
Morrone A *see* Kilinç *et al*
Mortensen NJMC *see* Wheeler JMD *et al*
Mortier G *see* Cross SJ *et al*
see Faivre L *et al*
Mortier G *et al*. Clinical and radiographic features of a family with hypochondroplasia owing to a novel Asn540Ser mutation in the fibroblast growth factor receptor 3 gene: *letter*, **220**
Mortier GR *et al*. Report of five novel and one recurrent *COL2A1* mutations with analysis of genotype-phenotype correlation in patients with a lethal type II collagen disorder, **263**
Mortimer P *see* Bell R *et al*
Mountford R *see* Pook MA *et al*
Mroczek K *see* Stankiewicz P *et al*
Mueller E *see* Zhou X-P *et al*
Mueller RF *see* Hutchin TP *et al*
see Nicholls AC *et al*
Muller PY *see* Zumsteg U *et al*
Muller D *see* Correa-Cerro LS *et al*
Mulligan LM *see* Hansford JR and Mulligan LM
see Onochie CI *et al*
Munnich A *see* Cormier-Daire V *et al*
see Cormier-Daire V *et al*
see Faivre L *et al*
see Sanlaville D *et al*
Munroe D *see* Catchpoole D *et al*
Murch SH *see* Aldred MA *et al*
Murday VA *see* Bell R *et al*
Murken J *see* Schmidt H *et al*
Murkin A *see* Forrest MS *et al*
Murphy T *see* Narain Y *et al*
Murray A *see* Youings SA *et al*
Murrell A *see* Catchpoole D *et al*
Mussaffi-Georgy H *see* Meeks M *et al*
Myers SM *see* Onochie CI *et al*

Nadal M *et al*. Hereditary neuropathy with liability to pressure palsies: two cases with a reciprocal translocation t(16;17)(q12;p11.2) interrupting the *PMP22* gene: *letter*, **396**
Nagamitsu S *see* Obata K *et al*
Nagase S *see* de Vivo I *et al*
Nájera C *see* Sánchez F *et al*
Nakagawa T *see* Usami S-I *et al*
Nakar O *see* Lev D *et al*
Namba A *see* Usami S-I *et al*
Nanba E *see* Yamamoto T *et al*
Narain Y *et al*. The *ACE* gene and Alzheimer's disease susceptibility, **695**
Narazaki O *see* Saitoh S *et al*
Narod SA *see* Liede A *et al*
see Metcalfe KA *et al*
Ndah BV *et al*. Prenatal detection of trisomy for the entire long arm of chromosome 7: *letter*, **551**
Neary W *see* Evans DGR *et al*
Negri T *see* Barresi R *et al*
Neitzel H *see* Weimer J *et al*
Nelson I *see* Reardon W *et al*
Neumann PE *see* Beresford RG *et al*
Newbury-Ecob R *see* Cross SJ *et al*
Newcombe R *see* Gray J *et al*
Newman B *et al*. Molecular diagnosis is important to confirm suspected pseudoachondroplasia: *letter*, **64**
Newman J *see* Thomson JR *et al*
Newton V *see* Evans DGR *et al*
Nicholls AC *et al*. Clinical phenotypes and molecular characterisation of three patients with Ehlers-Danlos syndrome type VII: *letter*, **e33**
Nichols WC *see* Thomson JR *et al*
Nicol RO *see* Chapman C *et al*
Nicolao P *see* Xiang F *et al*
Niebuhr E *see* Bugge M *et al*
Niermeijer MF *see* Lodder LN *et al*
Nieto AA *see* Dabora SL *et al*
Nieuwint AWM *et al*. Rapid detection of microdeletions using fluorescence in situ hybridisation (FISH) on buccal smears: *letter*, **e4**
Ninis VN *see* Kilinç *et al*
Ninomiya H *see* Yamamoto T *et al*
Nissen H *see* Thiart R *et al*
Noakes M *see* O'Callaghan FJK *et al*
Noeker M *see* Gohlke BC *et al*
Nolen MC *see* Sanlaville D *et al*
Nopola-Hemmi J *et al*. Two translocations of chromosome 15q associated with dyslexia, **771**
Norman P *see* Brain K *et al*
see Gray J *et al*
Norman A *see* Pook MA *et al*
North KN *see* Ellaway CJ *et al*
Northrup H *see* Astrinidis A *et al*
Nusier M *see* Hart TC *et al*
Nuytinck L *see* Mortier G *et al*
see Mortier GR *et al*
Nuytinck L *et al*. Glycine to tryptophan substitution in type I collagen in a patient with OI type III: a unique collagen mutation, **371**

Obata K *et al*. Mutation analysis of the methyl-CpG binding protein 2 gene (*MECP2*) in patients with Rett syndrome: *letter*, **608**
Obersztyn E *see* Stankiewicz P *et al*
O'Brien A *see* Humphreys L *et al*
O'Brien R *see* de Vivo I *et al*
O'Callaghan FJK *et al*. Renal angiomyolipomata and learning difficulty in tuberous sclerosis complex: *letter*, **156**
O'Dorisio MS *see* Frühwald MC *et al*
Oei P *see* Robertson SP *et al*
Oertel R *see* Voullaire L *et al*
Ogata T *see* Kirsch S *et al*
Ohno K *see* Yamamoto T *et al*
Ohta Y *see* Yamamoto T *et al*
Oiso N *see* Saitoh S *et al*
Oitmaa E *see* Teder M *et al*
Oley C *see* Nicholls AC *et al*
Olmer R *see* Willemsen R *et al*
Omori I *see* Obata K *et al*
Onochie CI *et al*. Characterisation of the human *GFRα-3* locus and investigation of the gene in Hirschsprung disease, **674**
Oostheim W *see* Waterham HR *et al*
Oostra BA *see* Willemsen R *et al*
O'Rawe A *see* Meeks M *et al*
Osborne JP *see* O'Callaghan FJK *et al*
Oshimura M *see* Engel JR *et al*
Ovens-Raeder A *see* Weimer J *et al*
Overhauser J *see* Johnson EI *et al*
see Maiwald R *et al*

Overhauser J *see* Perfumo C *et al*
Özkinay F *et al*. Genotype-phenotype correlation in three homozygotes and nine compound heterozygotes for the cystic fibrosis mutation 2183AA→G shows a severe phenotype: *letter*, **307**
Özme Ş *see* Dinçer P *et al*

Page DC *see* Ion A *et al*
Pal T *see* Liede A *et al*
Pang C-P *see* Lam C-W *et al*
Park K-Y *et al*. Desmin splice variants causing cardiac and skeletal myopathy, **851**
Park S-S *see* Potocki L *et al*
Parker MJ *see* Hutchin TP *et al*
Parker MJ *et al*. Genetics and deafness: what do families want?: *letter*, **e26**
Parsons E *see* Brain K *et al*
see Gray J *et al*
Parsons PG *see* Ellaway CJ *et al*
Parsons R *see* de Vivo I *et al*
Partington M *see* Fagan K *et al*
see Turner G and Partington M
Passmore AP *see* McIlroy SP *et al*
Patel MS *see* Blanck C *et al*
Paterson AD, Petronis A. Age and sex based genetic locus heterogeneity in type 1 diabetes, **186**
Patiño-García A *et al*. Analysis of the human tumour necrosis factor-alpha (TNFα) gene promoter polymorphisms in children with bone cancer: *letter*, **789**
Patton MA *see* Heathcote K *et al*
see Ion A *et al*
Patuzzo C *et al*. No linkage or association of the *IL-4Rα* gene Q576R mutation with atopic asthma in Italian families: *letter*, **382**
Pauciulo MW *see* Thomson JR *et al*
Payne AM *see* Bessant DAR *et al*
Peacock A *see* Thomson JR *et al*
Peacocke M *see* Çelebi JT *et al*
Pearson PL *see* Hes FJ *et al*
Pelet A *see* Bahuau M *et al*
Peltomäki P *see* Aktan-Collan K *et al*
see Roth S *et al*
Pena J *see* Özkinay F *et al*
Penault-Llorca F *see* Vaurs-Barrière C *et al*
Pentchev PG *see* Yamamoto T *et al*
Penttinen M *see* Cross SJ *et al*
Percesepe A *see* Desai DC *et al*
Perfumo C *et al*. The first three mosaic cri du chat syndrome patients with two rearranged cell lines: *letter*, **967**
Perrett C *see* Ellis D *et al*
Pescollderungg L *see* Patuzzo C *et al*
Petek E *see* Emberger W *et al*
see Kroisel PM *et al*
Peters DJM *see* Petrij F *et al*
Petit C *see* Masmoudi S *et al*
Petrij F *et al*. Diagnostic analysis of Rubinstein-Taybi syndrome: five cosmids should be used for microdeletion detection and low number of protein truncating mutations, **168**
Petronis A *see* Paterson AD and Petronis A
Petty EM *see* Martin DM *et al*
Phillimore HE *see* D'Cruz LG *et al*
Pieke-Dahl S *et al*. Genetic heterogeneity of Usher syndrome type II: localisation to chromosome 5q, **256**
Pierluigi M *see* Perfumo C *et al*
Pietilä T *see* Mattila KM *et al*
Pignatti PF *see* Patuzzo C *et al*
Pinson MP *see* Sanlaville D *et al*
Pinson M-P *see* Sidwell RU *et al*
Plant KE *et al*. Somatic mosaicism associated with a mild Alport syndrome phenotype: *letter*, **238**
Plaschke J *et al*. *BRCA2* germline mutations among early onset breast cancer patients unselected for family history of the disease: *letter*, **e17**
Plass C *see* Frühwald MC *et al*
see Zhou X-P *et al*
Plauchu H *see* Morlé L *et al*
Plecko-Startinig B *see* Emberger W *et al*
Polak-Charcon S *see* Meeks M *et al*
Poláková H *see* Zatková A *et al*
Pollitt RJ *see* Boyne J *et al*
Ponder BAJ *see* Seltmann M *et al*
Pook MA *et al*. Identification of three novel frameshift mutations in patients with Friedreich's ataxia: *letter*, **e38**
Poon PMK *see* Lam C-W *et al*
Pope FM *see* Nicholls AC *et al*
Porfirio B *see* Kilinç *et al*
Port RV *see* Chapman C *et al*
Porteous M *see* Campbell H *et al*
Porteous MEM *see* Berg JN *et al*
Portwine C *et al*. Absence of germline $p16^{INK4a}$ alterations in *p53* wild type Li-Fraumeni syndrome families: *letter*, **e13**
Potočnik U *see* Ravnik-Glavač M *et al*
Potocki L *et al*. Circadian rhythm abnormalities of melatonin in Smith-Magenis syndrome, **428**
Poulton KP *see* Barrett TG *et al*
Pound M *see* Youings SA *et al*
Poyhonen M. A clinical assessment of neurofibromatosis type 1 (NF1) and segmental NF in northern Finland: *letter*, **e43**
Poyhonen M *et al*. Epidemiology of neurofibromatosis type 1 (NF1) in northern Finland: *letter*, **632**
Pozzi S *see* Montera M *et al*
Praphanphoj V *et al*. Cryptic subtelomeric translocations in the 22q13 deletion syndrome, **58**
Precious S *see* Lynch SA *et al*
Preece MA *see* Wakeling EL *et al*
see Wakeling EL *et al*
Presneau N *see* Vaurs-Barrière C *et al*
Priestley MD *see* Green EK *et al*
Prieto F *see* Sánchez F *et al*
Prior TW *see* Zhou X-P *et al*
Priston M *see* Martin SN *et al*
Pritchard-Jones K *et al*. Familial Wilms tumour resulting from *WT1* mutation: intronic polymorphism causing artefactual constitutional homozygosity: *letter*, **377**
Pritchard M *see* Nadal M *et al*
Probst FJ *see* Martin DM *et al*
Protopapos A *see* Browne CE *et al*
Pulleyn LJ *see* Hutchin TP *et al*
Punnett HH *see* Johnson EI *et al*
Purcell K *see* Burton KA *et al*

Quaba AA *see* Berg JN *et al*
Quak E *see* Zschocke J *et al*
Quinn A *see* Moore SJ *et al*

Radice P *see* Desai DC *et al*
Ragnarsson G *see* Sigbjörnsdottir BI *et al*
Ragoussis J *see* Mirza G *et al*
Raguenes O *see* Chen J-M *et al*
Rahman N *see* McEntagart M *et al*
see Pritchard-Jones K *et al*
Ralston SH *see* Janssens K *et al*
Ramos J *see* Browne CE *et al*
Ramos-Arroyo MA *see* Nadal M *et al*
Ramsden RT *see* Evans DGR *et al*
Ranke MB *see* Mergenthaler S *et al*
Rappold GA *see* Gohlke BC *et al*
see Kirsch S *et al*
Rasmussen K *see* Bugge M *et al*
Rasmussen KR *see* Bugge M *et al*
Rasore-Quartino A *see* Wuyts W *et al*
Ravnik-Glavač M *et al*. Incidence of germline *hMLH1* and *hMSH2* mutations (HNPCC patients) among newly diagnosed colorectal cancers in a Slovenian population: *letter*, **533**
Raymond GV *see* Praphanphoj V *et al*
Reardon W *et al*. Evidence for digenic inheritance in some cases of Antley-Bixler syndrome?, **26**
Reddihough D *see* Halliday JL *et al*
Redmond EK *see* Davies JF *et al*
Reed JB *see* Hart PS *et al*
Rehal P *see* Hes FJ *et al*
Reid E *see* Baralle D *et al*
see de Lange RPJ *et al*
see Lindsey JC *et al*
Reik W *see* Catchpoole D *et al*
see Engel JR *et al*
Reiter JL *see* Frühwald MC *et al*
Reiter RJ *see* Potocki L *et al*
Renard J-P *see* Mortier G *et al*
Renieri A *see* Bruttini M *et al*
Resta N *see* Montera M *et al*
Rey-Berthod C *see* Hutter P *et al*
Reynolds PR *see* Dahia PM *et al*
Richard I *see* Dinçer P *et al*
Richard P *see* Forissier J-F *et al*
Richard S *see* Levy M and Richard S
Rickard S *et al*. Importance of clinical evaluation and molecular testing in the branchio-oto-renal (BOR) syndrome and overlapping phenotypes: *letter*, **623**
Riddell DC *see* Beresford RG *et al*
Riegel M *see* Kotzot D *et al*
see Röthlisberger B *et al*
Riess O *see* Wieczorek D *et al*
Rigardetto R *see* Silengo M and Rigardetto R
Riise R *see* Warburg M *et al*
Rimoin DL *see* Mortier GR *et al*
Rinne JO *see* Mattila KM *et al*
Ris-Stalpers C *see* Eussen BHJ *et al*
Rivard SR *et al*. Correlation between mutations and age in cystic fibrosis in a French Canadian population: *letter*, **225**
Rizzari G *see* Bruttini M *et al*
Roberts E *see* Deeble VJ *et al*
see Gray J *et al*
Roberts MH *see* Gao P-S *et al*

Robertson A *see* Voullaire L *et al*
Robertson SP *et al*. Homozygous deletion of *SHOX* in a mentally retarded male with Langer mesomelic dysplasia: *letter*, **959**
Robinson PN, Godfrey M. The molecular genetics of Marfan syndrome and related microfibrillopathies: *review*, **9**
Robinson WP *see* Kosaki K *et al*
Robson L *see* Koorey D *et al*
Rocchi M *see* Heller A *et al*
Rogers C *see* Brain K *et al*
see Gray J *et al*
Rogers P *see* Thomson JR *et al*
Romeijn GJ *see* Waterham HR *et al*
Romeo G *see* Costa M *et al*
Romio L *see* Montera M *et al*
Rosatelli MC *see* Iolascon A *et al*
Rosenblatt DS *see* Watkins D *et al*
Roth S *et al*. Mutation analysis of *SMAD2*, *SMAD3*, and *SMAD4* genes in hereditary non-polyposis colorectal cancer: *letter*, **298**
Röthlisberger B *see* Brecevic L *et al*
Röthlisberger B *et al*. A supernumerary marker chromosome originating from two different regions of chromosome 18, **121**
Röyttä M *see* Mattila KM *et al*
Rubinstein JH *see* Petrij F *et al*
Rubinsztein DC *see* Keramatipour M *et al*
see Lindsey JC *et al*
see Narain Y *et al*
Rudeberg A *see* Sobetzko D *et al*
Ruggieri M *see* John AM *et al*
Ruhenstroth E *see* Weimer J *et al*
Ruiz A *see* Borrego S *et al*
see Marcos I *et al*
Rumsby G *see* Reardon W *et al*
Rush LJ *see* Frühwald MC *et al*

Saad STO *see* Vasconcellos JPC *et al*
Saal HM *see* Petrij F *et al*
Saez ME *see* Borrego S *et al*
Saif R *see* Kelly TE *et al*
Sainio M *see* Evans DGR *et al*
Saitoh S *et al*. Oculocutaneous albinism type 2 with a *P* gene missense mutation in a patient with Angelman syndrome: *letter*, **392**
Sakati NA *see* Kelly TE *et al*
Sala E *see* Villa N *et al*
Salandi A *see* Barresi R *et al*
Salat U *et al*. Increase of FMRP expression, raised levels of *FMR1* mRNA, and clonal selection in proliferating cells with unmethylated fragile X repeat expansions: a clue to the sex bias in the transmission of full mutations?, **842**
Sampson J *see* Gray J *et al*
Sancak Ö *see* Dinçer P *et al*
Sancandi M *see* Costa M *et al*
Sánchez F *et al*. A constitutional homozygous mutation in the *RB1* gene in a patient with unilateral retinoblastoma: *letter*, **615**
Sandy CJ *see* Bell R *et al*
Sanjad SA *see* Kelly TE *et al*
Sanlaville D *et al*. Maternal uniparental heterodisomy of chromosome 14: chromosomal mechanism and clinical follow up, **525**
Santer R *see* Hillebrand G *et al*
Saraiva JM. Anticipation in progressive diaphyseal dysplasia: *letter*, **394**
Sargent CA *see* Blanco P *et al*
Sargent CA *et al*. Five cases of isolated glycerol kinase deficiency, including two families: failure to find genotype:phenotype correlation, **434**
Sarraf P *see* Zhou X-P *et al*
Sart DD *see* Massie J *et al*
Sasaki S *see* Gao P-S *et al*
Sato S *see* Kosaki K *et al*
Savarirayan R *see* Robertson SP *et al*
Savarirayan R *et al*. Rough skin, brittle hair, and photosensitivity: a mild phenotypic variant of trichothiodystrophy: *letter*, **312**
Savov *see* Kilinç *et al*
Scambler PJ *see* Robertson SP *et al*
Schackert HK *see* Plaschke J *et al*
Scharfe C *et al*. A novel mutation in the thiamine responsive megaloblastic anaemia gene *SLC19A2* in a patient with deficiency of respiratory chain complex I, **669**
Scheffer H *et al*. Two novel germline mutations of the retinoblastoma gene (*RB1*) that show incomplete penetrance, one splice site and one missense: *letter*, **e6**
Schinzel A *see* Brecevic L *et al*
see Fokstuen S and Schinzel A
see Kotzot D *et al*
see Röthlisberger B *et al*
Schinzel A *see* Faber J *et al*
Schmidt H *et al*. Mosaicism for a dup(12)(q22q13) in a patient with hypomelanosis of Ito and asymmetry: *letter*, **804**
Schofield PN *see* Catchpoole D *et al*
see Engel JR *et al*
Scholtz CL *see* Thiart R *et al*
Schuffenhauer S *see* Kotzot D *et al*
see Lichtner P *et al*
Schuffenhauer S *see* Schmidt H *et al*
Schwanitz G *see* Trost D *et al*
Schwartz CE *see* Cabezas DA *et al*
Schwartz K *see* Forissier J-F *et al*
Schwindt P *see* Trost D *et al*
Scott A *see* Metcalfe KA *et al*
Scott G *see* Ellis D *et al*
Scott-Brown M *see* Barrett TG *et al*
Seculi JL *see* Özkinay F *et al*
Seidel H *see* Kotzot D *et al*
Seidel J *see* Heller A *et al*
Seller A *see* Barrett TG *et al*
Seltmann M *et al*. A case of inv dup(8p) with early onset breast cancer: *letter*, **70**
Selvaag E. Pili torti et canaliculi and agenesis of the teeth: report of a new "pure" hair-tooth ectodermal dysplasia in a Norwegian family: *letter*, **721**
Semino-Mora C *see* Park K-Y *et al*
Sendtner M *see* Coovert DD *et al*
Senger G *see* Heller A *et al*
Seow WK *see* Hart PS *et al*
Seri M *see* Costa M *et al*
Seta N *see* Cormier-Daire V *et al*
see Vuillaumier-Barrot S *et al*
Setterfield B *see* Hastings RJ *et al*
Sevens C *see* Abramowicz MJ *et al*
Shaffer LG *see* Kosaki K *et al*
see Potocki L *et al*
Shapira L *see* Hart TC *et al*
Sharples PM *see* Stalker DJ *et al*
Sharrock A *see* Youings SA *et al*
Shaw DJ *see* de Lange RPJ *et al*
Shaw PJ *see* Lindsey JC *et al*
Shears DJ *see* Robertson SP *et al*
Shelling AN *see* Burton KA *et al*
Shendrik I *see* Çelebi JT *et al*
Sher JL *see* Nicholls AC *et al*
Shinkawa H *see* Abe S *et al*
see Usami S-I *et al*
Shirakawa T *see* Gao P-S *et al*
Shlumukova M *see* Blanco P *et al*
Siderovski DP *see* Bessant DAR *et al*
Sidransky E *see* Stone DL *et al*
Sidwell RU *et al*. Pure trisomy 20p resulting from isochromosome formation and whole arm translocation: *letter*, **454**
Siebert R *see* Hillebrand G *et al*
Sierrasesúmaga L *see* Patiño-Garcia A *et al*
Sigbjörnsdottir BI *et al*. Chromosome 8p alterations in sporadic and *BRCA2* 999del5 linked breast cancer, **342**
Silengo M, Rigardetto R. Hall-Riggs syndrome: a possible second affected family?: *letter*, **886**
Sillence D *see* Henderson S *et al*
see Reardon W *et al*
Silvers DN *see* Çelebi JT *et al*
Simard F *see* Rivard SR *et al*
Simard J *see* Badzioch M *et al*
Simeoni E *see* Hillebrand G *et al*
Simone C *see* Montera M *et al*
Simpson H *see* Meeks M *et al*
Sinclair RD *see* Savarirayan R *et al*
Slater H *see* Voullaire L *et al*
Slaugh R *see* Cabezas DA *et al*
Slavotinek A, Kingston H. Punctate calcification of the epiphyses, visceral malformations, and craniofacial dysmorphism in a female baby: *letter*, **796**
Slavotinek AM *et al*. Partial tetrasomy 21 in a male infant: *letter*, **e30**
Smallwood A *see* Engel JR *et al*
Smiraglia DJ *see* Frühwald MC *et al*
Smith A *see* Koorey D *et al*
see Reardon W *et al*
Smith SD *et al*. Tietz syndrome (hypopigmentation/deafness) caused by mutation of *MITF* , **446**
Smith WM *see* Zhou X-P *et al*
Smith M *see* Meiser B *et al*
Smolarek T *see* Astrinidis A *et al*
Snow BE *see* Bessant DAR *et al*
Sobetzko D *et al*. Achondroplasia with the *FGFR3* 1138g→a (G380R) mutation in two sibs sharing a 4p haplotype derived from their unaffected father: *letter*, **958**
Song J *see* Stratakis CA *et al*
Soskolne WA *see* Hart TC *et al*
Sotillo-Piñeiro E *see* Patiño-García A *et al*
Soubjaki V *see* Fagan K *et al*
Speicher MR *see* Schmidt H *et al*
Speizer FE *see* de Vivo I *et al*
Spiden S *see* Meeks M *et al*
Spranger S *see* Cross SJ *et al*
Spurlock G *see* McEntagart M *et al*
St Clair DM *see* de Lange RPJ *et al*

Stabholz A *see* Hart TC *et al*
Stalker DJ *et al*. Distal trisomy 2p and arachnodactyly: *letter*, **974**
Stanier P *see* Wakeling EL *et al*
see Wakeling EL *et al*
Stankiewicz P *et al*. Identification of supernumerary marker chromosomes derived from chromosomes 5, 6, 19, and 20 using FISH, **114**
Starke H *see* Heller A *et al*
see Stankiewicz P *et al*
Stead JA *see* Ndah BV *et al*
Steel CM *see* Hodgson SV *et al*
Steel CM *see* Young IE *et al*
Steinbach P *see* Salat U *et al*
Steinmann B *see* Faber J *et al*
Stevenson RE *see* Cabezas DA *et al*
Stone DL *et al*. Cardiovascular fibrosis, hydrocephalus, ophthalmoplegia, and visceral involvement in an American child with Gaucher disease: *letter*, **e40**
Stoppa-Lyonnet D *see* Hodgson SV *et al*
Storme L *see* Manouvrier-Hanu S *et al*
Stott NS *see* Chapman C *et al*
Stowasser M *see* Lafferty AR *et al*
Strachan T *see* Baser ME *et al*
see Lynch SA *et al*
Stratakis CA *see* Lafferty AR *et al*
Stratakis CA *et al*. Mapping of the human genes (*SLC23A2* and *SLC23A1*) coding for vitamin C transporters 1 and 2 (SVCT1 and SVCT2) to 5q23 and 20p12, respectively: *letter*, **e20**
Stratton MR *see* Mangion J *et al*
Superti-Furga A *see* Faber J *et al*
see Sobetzko D *et al*
Sutherland J *see* Martin SN *et al*
Suthers G *see* Meiser B *et al*
Svennevik EC *see* Hastings RJ *et al*
see Sidwell RU *et al*
Sykes B *see* Henderson S *et al*
Syngal S *et al*. Sensitivity and specificity of clinical criteria for hereditary non-polyposis colorectal cancer associated mutations in *MSH2* and *MLH1* , **641**
Syrris P *see* Heathcote K *et al*
Szudek J *et al*. Growth in North American white children with neurofibromatosis 1 (NF1), **933**

Taine L *see* Bahuau M *et al*
Taipale M *see* Nopola-Hemmi J *et al*
Takata A. Constitutional *WT1* mutations correlate with clinical features in children with progressive nephropathy: *letter*, **698**
Takeda K *see* Park K-Y *et al*
Tamura T *see* Johanning GL *et al*
Tan ASC *see* Goh DLM *et al*
Tan D-X *see* Potocki L *et al*
Tan E *see* Dinçer P *et al*
Tang *see* see Catchpoole D *et al*
Tanner MS *see* Burwinkel B *et al*
Tan J *see* Cormier-Daire V *et al*
Taphoorn MJB *see* Hes FJ *et al*
Tatlidil C *see* Beresford RG *et al*
Tayebi N *see* Stone DL *et al*
Taylor R *see* D'Cruz LG *et al*
Taylor CJ *see* Boyne J *et al*
Taymans SE *see* Lafferty AR *et al*
see Stratakis CA *et al*
Teare MD *see* Forrest MS *et al*
Tebbs S *see* Keramatipour M *et al*
Teder M *et al*. Distribution of *CFTR* gene mutations in cystic fibrosis patients from Estonia: *letter*, **e16**
Teebi AS. Hypoparathyroidism, retarded growth and development, and dysmorphism or Sanjad-Sakati syndrome: an Arab disease reminiscent of Kenny-Caffey syndrome: *letter*, **145**
Telvi L *see* Ion A *et al*
Temple IK *see* Marsh AJ *et al*
Tenenholz B *see* Johnson EI *et al*
Ten Kate LP *see* Nieuwint AWM *et al*
see Terhal PA *et al*
Tennstedt C *see* Knoblauch H *et al*
Tentler D *et al*. A microdeletion in 19q13.2 associated with mental retardation, skeletal malformations, and Diamond-Blackfan anaemia suggests a novel contiguous gene syndrome, **128**
Terhal PA *et al*. Breast hypoplasia and disproportionate short stature in the ear, patella, short stature syndrome: expansion of the phenotype?: *letter*, **719**
Testa F *see* Tiberio G *et al*
Thiart R *et al*. Predominance of a 6 bp deletion in exon 2 of the LDL receptor gene in Africans with familial hypercholesterolaemia, **514**
Thiel G *see* Knoblauch H *et al*
Thomas G *see* López-Correa C *et al*
Thomas GH *see* Praphanphoj V *et al*
Thomas NH *see* Pook MA *et al*
Thompson E *see* Meiser B *et al*
Thomson JR *et al*. Sporadic primary pulmonary hypertension is associated with germline mutations of the gene encoding BMPR-II, a receptor member of the TGF-β family, **741**
Tiberio G *et al*. Liebenberg syndrome: brachydactyly with joint dysplasia (MIM 186550): a second family: *letter*, **548**
Tigaud I *see* Cudry S *et al*
Tinschert S *see* Knoblauch H *et al*
Tolun A *see* Kilinç *et al*
Tomaras C *see* Koorey D *et al*
Tomiuk J *see* Mergenthaler S *et al*
Tomlinson IPM *see* Houlston RS and Tomlinson IPM
Tommerup N *see* Bugge M *et al*
Tong S-F *see* Lam C-W *et al*
Tönnies H *see* Weimer J *et al*
Tono T *see* Usami S-I *et al*
Topaloğlu H *see* Dinçer P *et al*
Torpy DJ *see* Lafferty AR *et al*
Trabetti E *see* Patuzzo C *et al*
Trembath RC *see* Aldred MA *et al*
see Thomson JR *et al*
Trijsburg RW *see* Lodder LN *et al*
Trompeter R *see* Rickard S *et al*
Trost D *et al*. Investigation of meiotic rearrangements in DGS/VCFS patients with a microdeletion 22q11.2: *letter*, **452**
Tsukumo DML *see* Vasconcellos JPC *et al*
Tsutsumi Y *see* Yamamoto T *et al*
Tucker K *see* Meiser B *et al*
Tukel T *see* Nuytinck L *et al*
Turner G, Partington M. Recurrence risks in undiagnosed mental retardation: *letter*, **e45**
Turner G *see* Fagan K *et al*
Turnpenny P *see* Moore SJ *et al*
Turnpenny PD, Hole R. The first description of lethal pterygium syndrome with facial clefting (Bartsocas-Papas syndrome) in 1600: *letter*, **314**

Uhlhaas S *see* Trost D *et al*
UK Huntington's Disease Prediction Consortium *see* Harper PS *et al*
Uhrig S *see* Schmidt H *et al*
Upadhyaya M *see* John AM *et al*
Urtizberea JA *see* Dinçer P *et al*
Usami S-I *see* Abe S *et al*
Usami S-I *et al*. Prevalence of mitochondrial gene mutations among hearing impaired patients, **38**
Uutela A *see* Aktan-Collan K *et al*
Uygur C *see* Hart PS *et al*

Vahidassr MD *see* McIlroy SP *et al*
Valiente A *see* Nadal M *et al*
van Bever Y *see* Terhal PA *et al*
van Broeckhoven C *see* van Goethem G *et al*
Vandenberghe A *see* Morlé L *et al*
van den Berghe JA *see* Goh DLM *et al*
van den Ouweland AMW *see* Eussen BHJ *et al*
van den Ouweland A *see* Dabora SL *et al*
van der Burgt I *see* Ion A *et al*
van der Luijt RB *see* Hes FJ *et al*
van der Smagt JJ *see* Hes FJ *et al*
see Petrij F *et al*
van der Vlies P *see* Scheffer H *et al*
van Dyke TE *see* Hart TC *et al*
van Ee CC *see* Burton KA *et al*
van Esch H *see* Lichtner P *et al*
van Goethem G *et al*. Further evidence for genetic heterogeneity of autosomal dominant disorders with accumulation of multiple deletions of mitochondrial DNA: *letter*, **547**
van Haeringen A *see* Petrij F *et al*
van Hagen JM *see* Nieuwint AWM *et al*
van Hemel JO *see* Eussen BHJ *et al*
Vanhoenacker F *see* Janssens K *et al*
see Wuyts W *et al*
Vanhorne JB *see* Onochie CI *et al*
van Hul E *see* Janssens K *et al*
van Hul W *see* Janssens K *et al*
see Wuyts W *et al*
Vanier MT *see* Yamamoto T *et al*
van Karnebeek CD *see* Petrij F *et al*
van Maldergem L *see* Janssens K *et al*
van Ommen G-JB *see* Petrij F *et al*
van Ravenswaaij C *see* Cross SJ *et al*
van Reen M *see* Ion A *et al*
Variend D *see* Pritchard-Jones K *et al*
Varnava A *see* D'Cruz LG *et al*
Vasconcellos JPC *et al*. Novel mutation in the *MYOC* gene in primary open angle glaucoma patients: *letter*, **301**
Vasen H *see* Hodgson SV *et al*
Vaurs-Barrière C *et al*. Low frequency of microsatellite instability in *BRCA1* mutated breast tumours: *letter*, **e32**
Vaux C *see* Voullaire L *et al*
Vekemans M *see* Sanlaville D *et al*
Vellodi A *see* Beesley CE *et al*

Verbruggen LA *see* Janssens K *et al*
Verellen-Dumoulin C *see* Chen J-M *et al*
Vergnes P *see* Bahuau M *et al*
Vergotine J *see* Thiart R *et al*
Verhoef S *see* Eussen BHJ *et al*
Verlind E *see* Scheffer H *et al*
Vermaak WJH *see* Thiart R *et al*
Verselis S *see* Portwine C *et al*
Vetrie D *see* Plant KE *et al*
Vidaud D *see* Bahuau M *et al*
Vidaud M *see* Bahuau M *et al*
Vigneswaren S *see* Stalker DJ *et al*
Villa N *et al*. Monosomy and trisomy 1q44-qter in two sisters originating from a half cryptic 1q;15p translocation: *letter*, **612**
Vitali A *see* Barresi R *et al*
Vitelli F *see* Bruttini M *et al*
Vogel W *see* Correa-Cerro LS *et al*
see Jakubowski L *et al*
Volpe MD *see* Bruttini M *et al*
von Deimling A *see* Zhou X-P *et al*
Voullaire L *et al*. Chromosomal duplication of band 10p14 segregating through four generations: *letter*, **233**
Voutilainen A *see* Nopola-Hemmi J *et al*
Vuillaumier-Barrot S *see* Cormier-Daire V *et al*
Vuillaumier-Barrot S *et al*. Identification of four novel *PMM2* mutations in congenital disorders of glycosylation (CDG) Ia French patients, **579**

Wada T *see* Saitoh S *et al*
Wagner K *see* Emberger W *et al*
see Kroisel PM *et al*
Wakeling EL *et al*. Biallelic expression of *IGFBP1* and *IGFBP3*, two candidate genes for the Silver-Russell syndrome: *letter*, **65**
Silver-Russell syndrome and ring chromosome 7: *letter*, **380**
Waldron HA. A case of dyschondrosteosis from Roman Britain: *letter*, **e27**
Walker S *see* Hart TC *et al*
Walker-Smith JA *see* Aldred MA *et al*
Wallace A *see* Evans DGR *et al*
Wallace AJ *see* Baser ME *et al*
Wallerstein R *see* Petrij F *et al*
Wallis GA *see* Gregory CA *et al*
Walne A *see* Meeks M *et al*
Wanders RJA *see* Waterham HR *et al*
Wang J *et al*. A novel mutation in the *CFTR* gene correlates with severe clinical phenotype in seven Hispanic patients: *letter*, **215**
Wang Y *see* Lynch SA *et al*
Warburg M *et al*. Bardet-Biedl and Cohen syndromes: differential diagnostic criteria: *letter*, **e46**
Ward K *see* Thomson JR *et al*
Watemberg N *see* Lev D *et al*
Waterham HR *et al*. Incidence and molecular mechanism of aberrant splicing owing to a G→C splice acceptor site mutation causing Smith-Lemli-Opitz syndrome: *letter*, **387**
Waters J *see* Green EK *et al*
Watkins D *et al*. Complementation studies in the *cblA* class of inborn error of cobalamin metabolism: evidence for interallelic complementation and for a new complementation class (*cblH*), **510**
Watts S *see* Ellis D *et al*
Webb T *see* Hampson K *et al*
Webster AR *et al*. Risk of multisystem disease in isolated ocular angioma (haemangioblastoma): *letter*, **62**
Weimer J *et al*. Delineation of a complex karyotypic rearrangement by microdissection and CGH in a family affected with split foot, **442**
Weintraub S *see* Lev D *et al*
Weis M *see* Mortier GR *et al*
Weiss B *see* Kirsch S *et al*
Welch JP *see* Beresford RG *et al*
Wellesley D *see* Marsh AJ *et al*
Wells D *see* Hastings RJ *et al*
Wenger G *see* Frühwald MC *et al*
Wenger SL *see* Ndah BV *et al*
Wenstrom KD *see* Johanning GL *et al*
Wertz K *see* Wang J *et al*
Westman J *et al*. Efficacy of a touchscreen computer based family cancer history questionnaire and subsequent cancer risk assessment, **354**
Wheeler JMD *et al*. The role of hypermethylation of the *hMLH1* promoter region in HNPCC versus MSI+ sporadic colorectal cancers, **588**
Wheeler L *see* Thomson JR *et al*
White KD *see* Lindsey JC *et al*
White PS *see* Frühwald MC *et al*
Whiteford ML *et al*. A child with bisatellited, dicentric chromosome 15 arising from a maternal paracentric inversion of chromosome 15q: *letter*, **e11**
Whittaker J *see* Hes FJ *et al*
Whittall RA *see* Heath KE *et al*
Wiebe W *see* Trost D *et al*
Wieczorek D *et al*. Unexpected high frequency of de novo unbalanced translocations in patients with Wolf-Hirschhorn syndrome (WHS): *letter*, **798**
Wiedemann U *see* Weimer J *et al*
Wilkie AOM *see* Lynch SA *et al*
Wilkin DJ *see* Mortier GR *et al*
Willatt LR *see* Seltmann M *et al*
Willemsen R *et al*. Twin sisters, monozygotic with the fragile X mutation, but with a different phenotype, **603**
Williams D *see* Hart PS *et al*
Williamson R *see* Delatycki MB *et al*
Wilson DI *see* Jones EA *et al*
Winchester BG *see* Beesley CE *et al*
Winship I *see* Burton KA *et al*
Winter RM *see* Reardon W *et al*
see Robertson SP *et al*
Winterpacht A *see* Faber J *et al*
Wirth J *see* Heller A *et al*
Wöhrle D *see* Salat U *et al*
Wollmann HA *see* Mergenthaler S *et al*
Wong L-JC *see* Wang J *et al*
Wong V *see* Lam C-W *et al*
Woods CG *see* Deeble VJ *et al*
see Hampson K *et al*
Woodward ER *et al*. Familial clear cell renal cell carcinoma (FCRC): clinical features and mutation analysis of the *VHL*, *MET*, and *CUL2* candidate genes, **348**
Wouters C *see* de Ravel TJL *et al*
Wright FA *see* Borrego S *et al*
see Desai DC *et al*
Wright JT *see* Hart PS *et al*
Wright MJ *et al*. Achondroplasia and nail-patella syndrome: the compound phenotype: *letter*, **e25**
Wright MJ *see* Nicholls AC *et al*
Wuyts W *et al*. The *ALX4* homeobox gene is mutated in patients with ossification defects of the skull (foramina parietalia permagna, OMIM 168500), **916**
Wyllie AH *see* Young IE *et al*

Xiang F *et al*. Mutation screening in Rett syndrome patients, **250**
Xuereb J *see* Narain Y *et al*

Yacoub M *see* Thomson JR *et al*
Yamakawa K *see* Yamamoto T *et al*
Yamamoto T *et al*. Genotype-phenotype relationship of Niemann-Pick disease type C: a possible correlation between clinical onset and levels of NPC1 protein in isolated skin fibroblasts: *letter*, **707**
Yamashita Y *see* Obata K *et al*
Yanoov M *see* Lev D *et al*
Yassin OM *see* Hart TC *et al*
Yates AJ *see* Zhou X-P *et al*
Yates JRW, Moore AT. Genetic susceptibility to age related macular degeneration: *review*, **83**
Yates JRW *see* Leggatt V *et al*
Yeung W-L *see* Lam C-W *et al*
Yip A *see* Narain Y *et al*
Youings SA *et al*. FRAXA and FRAXE: the results of a five year survey, **415**
Young EP *see* Beesley CE *et al*
Young ID *see* Hutchin TP *et al*
see Parker MJ *et al*
Young IE *et al*. The CAG repeat within the androgen receptor gene in male breast cancer patients: *letter*, **139**
Yuen ST *see* Desai DC *et al*

Zabel B *see* Faber J *et al*
see Gregory CA *et al*
Zammarchi E *see* Kilinç *et al*
Zanoni G *see* Patuzzo C *et al*
Zara F *see* Perfumo C *et al*
Zatková A *et al*. Novel mutations in the homogentisate-1,2-dioxygenase gene identified in Slovak patients with alkaptonuria: *letter*, **539**
Zewald RA *see* Hes FJ *et al*
Zhang Y *see* Hart PS *et al*
see Hart TC *et al*
see Hart TC *et al*
Zhang Z *see* Xiang F *et al*
Zhou X-P *et al*. Over-representation of *PPARγ* sequence variants in sporadic cases of glioblastoma multiforme: preliminary evidence for common low penetrance modifiers for brain tumour risk in the general population, **410**
Zierler H *see* Emberger W *et al*
Zimak A *see* Humphreys L *et al*
Zlorogorski A *see* Hart TC *et al*
Zschocke J *et al*. Mutation analysis in glutaric aciduria type I, **177**
Zucman-Rossi J *see* López-Correa C *et al*
Zudik A *see* Lev D *et al*

Zumsteg U *et al.* Alström syndrome: confirmation of linkage to chromosome 2p12-13 and phenotypic heterogeneity in three affected sibs: *letter*, **e8**

Zvarík M *see* Zatková A *et al*

VOLUME 37 • SUBJECT INDEX

18q− syndrome, No evidence for imprinting in distal 18q: *letter,* **152**
19q, A locus for primary ciliary dyskinesia maps to chromosome 19q, **241**

2157delG mutation, 2157delG: a frequent mutation in *BRCA2* missed by PTT: *letter,* **e42**
2183AA→G, Genotype-phenotype correlation in three homozygotes and nine compound heterozygotes for the cystic fibrosis mutation 2183AA→G shows a severe phenotype: *letter,* **307**
22qter deletion syndrome, Cryptic subtelomeric translocations in the 22q13 deletion syndrome, **58**

4p16.3 aneuploidy syndromes, Fine molecular mapping of the 4p16.3 aneuploidy syndromes in four translocation families: *letter,* **449**

5p deletion, The first three mosaic cri du chat syndrome patients with two rearranged cell lines: *letter,* **967**
5p duplication, The first three mosaic cri du chat syndrome patients with two rearranged cell lines: *letter,* **967**

abnormal phenotypes, Disease associated balanced chromosome rearrangements: a resource for large scale genotype-phenotype delineation in man, **858**
absent patella, Breast hypoplasia and disproportionate short stature in the ear, patella, short stature syndrome: expansion of the phenotype?: *letter,* **719**
***ACE* gene,** The *ACE* gene and Alzheimer's disease susceptibility, **695**
***ACE* I allele,** The *ACE* I allele is associated with increased risk for ruptured intracranial aneurysms, **498**
achondrogenesis II-hypochondrogenesis, Report of five novel and one recurrent *COL2A1* mutations with analysis of genotype-phenotype correlation in patients with a lethal type II collagen disorder, **263**
achondroplasia, Achondroplasia and nail-patella syndrome: the compound phenotype: *letter,* **e25**
Achondroplasia with the *FGFR3* 1138g→a (G380R) mutation in two sibs sharing a 4p haplotype derived from their unaffected father: *letter,* **958**
Germline and somatic mosaicism in achondroplasia: *letter,* **956**
acromesomelic dysplasia Maroteaux type, Exclusion of chromosome 9 helps to identify mild variants of acromesomelic dysplasia Maroteaux type, **52**
acromesomelic dysplasias, Exclusion of chromosome 9 helps to identify mild variants of acromesomelic dysplasia Maroteaux type, **52**
adenomatous polyposis coli, Deletion and duplication of the adenomatous polyposis coli gene resulting from an interchromosomal insertion involving 5(q22q23.3) in the father: *letter,* **141**
aetiology, Genetics and deafness: what do families want?: *letter,* **e26**
age of diagnosis, Age and sex based genetic locus heterogeneity in type 1 diabetes, **186**
age related macular degeneration, Genetic susceptibility to age related macular degeneration: *review,* **83**
Alagille syndrome, *JAGGED1* expression in human embryos: correlation with the Alagille syndrome phenotype, **658**
Albright hereditary osteodystrophy, Germline mosaicism for a *GNAS1* mutation and Albright hereditary osteodystrophy: *letter,* **e35**
aldosterone, A novel genetic locus for low renin hypertension: familial hyperaldosteronism type II maps to chromosome 7 (7p22), **831**
alkaptonuria, Alkaptonuria in Italy: polymorphic haplotype background, mutational profile, and description of four novel mutations in the homogentisate 1,2-dioxygenase gene: *letter,* **309**
Novel mutations in the homogentisate-1,2-dioxygenase gene identified in Slovak patients with alkaptonuria: *letter,* **539**
allelic mutations, Haim-Munk syndrome and Papillon-Lefèvre syndrome are allelic mutations in cathepsin C, **88**
Alport syndrome, Mosaicism in Alport syndrome and genetic counselling: *letter,* **717**
Somatic mosaicism associated with a mild Alport syndrome phenotype: *letter,* **238**
Alström syndrome, Alström syndrome: confirmation of linkage to chromosome 2p12-13 and phenotypic heterogeneity in three affected sibs: *letter,* **e8**
The continuing failure to recognise Alström syndrome and further evidence of genetic homogeneity: *letter,* **219**
ALX4, The *ALX4* homeobox gene is mutated in patients with ossification defects of the skull (foramina parietalia permagna, OMIM 168500), **916**
Alzheimer's disease, Butyrylcholinesterase K variant is genetically associated with late onset Alzheimer's disease in Northern Ireland, **182**
Dipeptidyl carboxypeptidase 1 (*DCP1*) and butyrylcholinesterase (*BCHE*) gene interactions with the apolipoprotein E ε4 allele as risk factors in Alzheimer's disease and in Parkinson's disease with coexisting Alzheimer pathology, **766**
The *ACE* gene and Alzheimer's disease susceptibility, **695**
anal atresia, Proximal 10q trisomy: a new case with anal atresia: *letter,* **e24**
androgen receptor gene, The CAG repeat within the androgen receptor gene in male breast cancer patients: *letter,* **139**
Angelman syndrome, Oculocutaneous albinism type 2 with a *P* gene missense mutation in a patient with Angelman syndrome: *letter,* **392**
anorectal malformation, Autosomal dominant sacral agenesis: Currarino syndrome: *review,* **561**
anticipation, Anticipation in progressive diaphyseal dysplasia: *letter,* **394**
Antley-Bixler syndrome, Evidence for digenic inheritance in some cases of Antley-Bixler syndrome?, **26**
anxiety, The psychological impact of a cancer family history questionnaire completed in general practice: *letter,* **470**
APC, Appendiceal carcinoma complicating adenomatous polyposis in a young woman with a de novo constitutional reciprocal translocation t(5;8)(q22;p23.1): *letter,* **71**
APOE, Butyrylcholinesterase K variant is genetically associated with late onset Alzheimer's disease in Northern Ireland, **182**
apolipoprotein B, Predominance of a 6 bp deletion in exon 2 of the LDL receptor gene in Africans with familial hypercholesterolaemia, **514**
appendix, Appendiceal carcinoma complicating adenomatous polyposis in a young woman with a de novo constitutional reciprocal translocation t(5;8)(q22;p23.1): *letter,* **71**
arachnodactyly, Distal trisomy 2p and arachnodactyly: *letter,* **974**
arteriohepatic dysplasia, *JAGGED1* expression in human embryos: correlation with the Alagille syndrome phenotype, **658**
asymmetry, Mosaicism for a dup(12)(q22q13) in a patient with hypomelanosis of Ito and asymmetry: *letter,* **804**
atopic asthma, Variants of *STAT6* (signal transducer and activator of transcription 6) in atopic asthma: *letter,* **380**
atopy, No linkage or association of the *IL-4Rα* gene Q576R mutation with atopic asthma in Italian families: *letter,* **382**
attitudes, Attitudes to genetic testing for breast cancer susceptibility in women at increased risk of developing hereditary breast cancer: *letter,* **472**
audiology, Use of MRI and audiological tests in presymptomatic diagnosis of type 2 neurofibromatosis (NF2): *letter,* **944**
audit, Evidence based medicine in practice: lessons from a Scottish clinical genetics project, **684**
autosomal dominant disorders, Further evidence for genetic heterogeneity of autosomal dominant disorders with accumulation of multiple deletions of mitochondrial DNA: *letter,* **547**
autosomal dominant hearing loss, A novel C202F mutation in the connexin26 gene (*GJB2*) associated with autosomal dominant isolated hearing loss, **368**
autosomal dominant inheritance, A family with hereditary port wine stain: *letter,* **e12**
The gene for branchio-oculo-facial syndrome does not colocalise to the *EYA1-4* genes: *letter,* **620**
AZFa, Divergent outcomes of intrachromosomal recombination on the human Y chromosome: male infertility and recurrent polymorphism, **752**

Bannayan-Riley-Ruvalcaba syndrome, Absence of germline mutations in *MINPP1,* a phosphatase encoding gene centromeric of *PTEN,* in patients with Cowden and Bannayan-Riley-Ruvalcaba syndrome without germline *PTEN* mutations: *letter,* **715**
Bardet-Biedl syndrome, Bardet-Biedl and Cohen syndromes: differential diagnostic criteria: *letter,* **e46**
Bartsocas-Papas syndrome, The first description of lethal pterygium syndrome with facial clefting (Bartsocas-Papas syndrome) in 1600: *letter,* **314**
BCHE K, Butyrylcholinesterase K variant is genetically associated with late onset Alzheimer's disease in Northern Ireland, **182**
Bcl-1, Does the survival motor neuron protein (SMN) interact with Bcl-2?: *letter,* **536**
Beckwith-Wiedemann syndrome, Epigenotype-phenotype correlations in Beckwith-Wiedemann syndrome, **921**
Mutation analysis of *H19* and *NAP1L4* (*hNAP2*) candidate genes and *IGF2* DMR2 in Beckwith-Wiedemann syndrome: *letter,* **212**
benign familial infantile convulsions, Benign familial infantile convulsions: report of a UK family and confirmation of genetic heterogeneity: *letter,* **e31**
birth defects, A clinical study of 57 children with fetal anticonvulsant syndromes, **489**

bisatellited dicentric chromosome 15, A child with bisatellited, dicentric chromosome 15 arising from a maternal paracentric inversion of chromosome 15q: *letter,* **e11**
***BMPR2*,** Sporadic primary pulmonary hypertension is associated with germline mutations of the gene encoding BMPR-II, a receptor member of the TGF-β family, **741**
BMPR-II, Sporadic primary pulmonary hypertension is associated with germline mutations of the gene encoding BMPR-II, a receptor member of the TGF-β family, **741**
BO, Importance of clinical evaluation and molecular testing in the branchio-oto-renal (BOR) syndrome and overlapping phenotypes: *letter,* **623**
BOF, The gene for branchio-oculo-facial syndrome does not colocalise to the *EYA1-4* genes: *letter,* **620**
bone cancer, Analysis of the human tumour necrosis factor-alpha (TNFα) gene promoter polymorphisms in children with bone cancer: *letter,* **789**
BOR, Importance of clinical evaluation and molecular testing in the branchio-oto-renal (BOR) syndrome and overlapping phenotypes: *letter,* **623**
The gene for branchio-oculo-facial syndrome does not colocalise to the *EYA1-4* genes: *letter,* **620**
brachydactyly, Liebenberg syndrome: brachydactyly with joint dysplasia (MIM 186550): a second family: *letter,* **548**
brachydactyly A1, Clinical and radiological assessment of a family with mild brachydactyly type A1: the usefulness of metacarpophalangeal profiles, **292**
brain tumour, Suggestive evidence for a site specific prostate cancer gene on chromosome 1p36: *letter,* **947**
branchio-oto-renal syndrome, A *SALL1* mutation causes a branchio-oto-renal syndrome-like phenotype: *letter,* **458**
***BRCA1* mutations,** Low prevalence of germline *BRCA1* mutations in early onset breast cancer without a family history: *letter,* **792**
***BRCA1*,** An evaluation of needs of female *BRCA1* and *BRCA2* carriers undergoing genetic counselling, **866**
Attitudes towards termination of pregnancy in subjects who underwent presymptomatic testing for the *BRCA1*/*BRCA2* gene mutation in The Netherlands: *letter,* **883**
Low frequency of microsatellite instability in *BRCA1* mutated breast tumours: *letter,* **e32**
***BRCA2*,** 2157delG: a frequent mutation in *BRCA2* missed by PTT: *letter,* **e42**
An evaluation of needs of female *BRCA1* and *BRCA2* carriers undergoing genetic counselling, **866**
Attitudes towards termination of pregnancy in subjects who underwent presymptomatic testing for the *BRCA1*/*BRCA2* gene mutation in The Netherlands: *letter,* **883**
Chromosome 8p alterations in sporadic and *BRCA2* 999del5 linked breast cancer, **342**
BRCA2 germline mutations among early onset breast cancer patients unselected for family history of the disease: *letter,* **e17**
breast cancer, A model protocol evaluating the introduction of genetic assessment for women with a family history of breast cancer, **192**
Can hair be used to screen for breast cancer?: *letter,* **297**
Chromosome 8p alterations in sporadic and *BRCA2* 999del5 linked breast cancer, **342**
Delineation of a new syndrome: clustering of pyloric stenosis, endometriosis, and breast cancer in two families: *letter,* **794**
Guidelines for a genetic risk based approach to advising women with a family history of breast cancer, **203**
Low frequency of microsatellite instability in *BRCA1* mutated breast tumours: *letter,* **e32**
BRCA2 germline mutations among early onset breast cancer patients unselected for family history of the disease: *letter,* **e17**
breast cancer genes, Clinical geneticists' attitudes and practice towards testing for breast cancer susceptibility genes: *letter,* **157**
breast hypoplasia, Breast hypoplasia and disproportionate short stature in the ear, patella, short stature syndrome: expansion of the phenotype?: *letter,* **719**
buccal smears, Rapid detection of microdeletions using fluorescence in situ hybridisation (FISH) on buccal smears: *letter,* **e4**
bullous skin lesion, Skin pigmentary anomalies in a mosaic form of partial tetrasomy 3q: *letter,* **723**
butyrylcholinesterase, Dipeptidyl carboxypeptidase 1 (*DCP1*) and butyrylcholinesterase (*BCHE*) gene interactions with the apolipoprotein E ε4 allele as risk factors in Alzheimer's disease and in Parkinson's disease with coexisting Alzheimer pathology, **766**

C202F mutation, A novel C202F mutation in the connexin26 gene (*GJB2*) associated with autosomal dominant isolated hearing loss, **368**
CADASIL, Screening British CADASIL families for mutations in the *NOTCH3* gene: *letter,* **224**
Camurati-Engelmann disease, Anticipation in progressive diaphyseal dysplasia: *letter,* **394**
Localisation of the gene causing diaphyseal dysplasia Camurati-Engelmann to chromosome 19q13, **245**
cancer, Detecting low penetrance genes in cancer: the way ahead: *review,* **161**
The psychological impact of a cancer family history questionnaire completed in general practice: *letter,* **470**
cancer genetics, An evaluation of needs of female *BRCA1* and *BRCA2* carriers undergoing genetic counselling, **866**
candidate gene, Mutation screening in Rett syndrome patients, **250**
candidate genes, Detailed mapping, mutation analysis, and intragenic polymorphism identification in candidate Noonan syndrome genes *MYL2*, *DCN*, *EPS8*, and *RPL6* : *letter,* **884**
capillary haemangiomata, Proximal 10q trisomy: a new case with anal atresia: *letter,* **e24**
carcinoma, Appendiceal carcinoma complicating adenomatous polyposis in a young woman with a de novo constitutional reciprocal translocation t(5;8)(q22;p23.1): *letter,* **71**
cardiac and skeletal myopathy, Desmin splice variants causing cardiac and skeletal myopathy, **851**
cardiac troponin T gene, Cytosine methylation confers instability on the cardiac troponin T gene in hypertrophic cardiomyopathy: *letter,* **e18**
cardiomyopathy, Disruption of heart sarcoglycan complex and severe cardiomyopathy caused by β sarcoglycan mutations, **102**
cardiovascular fibrosis, Cardiovascular fibrosis, hydrocephalus, ophthalmoplegia, and visceral involvement in an American child with Gaucher disease: *letter,* **e40**
care pathways, Evidence based medicine in practice: lessons from a Scottish clinical genetics project, **684**
carpal synostosis, Liebenberg syndrome: brachydactyly with joint dysplasia (MIM 186550): a second family: *letter,* **548**
carrier rates, Determination of the frequency of connexin26 mutations in inherited sensorineural deafness and carrier rates in the Tunisian population using DGGE: *letter,* **e39**
cataract, The genetics of childhood cataract: *review,* **481**
cathepsin C, Identification of cathepsin C mutations in ethnically diverse Papillon-Lefèvre syndrome patients, **927**
Localisation of a gene for prepubertal periodontitis to chromosome 11q14 and identification of a cathepsin C gene mutation, **95**
cathepsin C mutation, Haim-Munk syndrome and Papillon-Lefèvre syndrome are allelic mutations in cathepsin C, **88**
cationic trypsinogen gene, A CGC>CAT gene conversion-like event resulting in the R122H mutation in the cationic trypsinogen gene and its implication in the genotyping of pancreatitis: *letter,* **e36**
***cblA*,** Complementation studies in the *cblA* class of inborn error of cobalamin metabolism: evidence for interallelic complementation and for a new complementation class (*cblH*), **510**
***cblH*,** Complementation studies in the *cblA* class of inborn error of cobalamin metabolism: evidence for interallelic complementation and for a new complementation class (*cblH*), **510**
CDG, Identification of four novel *PMM2* mutations in congenital disorders of glycosylation (CDG) Ia French patients, **579**
***CDKN2A*,** Identification of *PTEN* mutations in metastatic melanoma specimens, **653**
centromere misdivision, Pure trisomy 20p resulting from isochromosome formation and whole arm translocation: *letter,* **454**
cerebral palsy, Hemiplegic cerebral palsy and the factor V Leiden mutation: *letter,* **787**
***CFTR* gene,** A novel mutation in the *CFTR* gene correlates with severe clinical phenotype in seven Hispanic patients: *letter,* **215**
***CFTR* gene mutations,** Distribution of *CFTR* gene mutations in cystic fibrosis patients from Estonia: *letter,* **e16**
***CFTR* mutation,** Correlation between mutations and age in cystic fibrosis in a French Canadian population: *letter,* **225**
***CFTR* mutations,** Many ΔF508 heterozygote neonates with transient hypertrypsinaemia have a second, mild *CFTR* mutation: *letter,* **543**
CGH, An analphoid supernumerary marker chromosome derived from chromosome 3 ascertained in a fetus with multiple malformations: *letter,* **807**
Delineation of a complex karyotypic rearrangement by microdissection and CGH in a family affected with split foot, **442**
Molecular cytogenetic characterisation of partial trisomy 9q in a case with pyloric stenosis and a review, **529**
channelopathies, Channelopathies: ion channel defects linked to heritable clinical disorders: *review,* **729**
Charcot-Marie-Tooth disease, Distal spinal muscular atrophy with vocal cord paralysis (dSMA-VII) is not linked to the *MPD2* locus on chromosome 5q31: *letter,* **e14**
CHARGE association, CHARGE association in a child with de novo chromosomal aberration 46,X,der(X)t(X;2)(p22.1;q33) detected by spectral karyotyping: *letter,* **e47**
cherubism, Familial craniofacial fibrous dysplasia: absence of linkage to *GNAS1* and the gene for cherubism: *letter,* **e37**
children, Genetics and deafness: what do families want?: *letter,* **e26**
choanal atresia, CHARGE association in a child with de novo chromosomal aberration 46,X,der(X)t(X;2)(p22.1;q33) detected by spectral karyotyping: *letter,* **e47**
cholesterol metabolism, The Smith-Lemli-Opitz syndrome: *review,* **321**
chromatin modification, Chromatin modification and disease: *review,* **905**
chromosome 10, *RET* genotypes comprising specific haplotypes of polymorphic variants predispose to isolated Hirschsprung disease, **572**

chromosome 10p14, Chromosomal duplication of band 10p14 segregating through four generations: *letter,* **233**
chromosome 12, Mosaicism for a dup(12)(q22q13) in a patient with hypomelanosis of Ito and asymmetry: *letter,* **804**
chromosome 13q14, Tandem triplication of chromosome 13q14 with inverted interstitial segment in a 4 year old girl: *letter,* **964**
chromosome 14, Maternal uniparental heterodisomy of chromosome 14: chromosomal mechanism and clinical follow up, **525**
chromosome 15, Two translocations of chromosome 15q associated with dyslexia, **771**
chromosome 15q, A child with bisatellited, dicentric chromosome 15 arising from a maternal paracentric inversion of chromosome 15q: *letter,* **e11**
chromosome 17, Interstitial deletion of chromosome 17 (del(17)(q22q23.3 confirms a link with oesophageal atresia: *letter,* **701**
chromosome 18, A supernumerary marker chromosome originating from two different regions of chromosome 18, **121**
chromosome 19q13, A microdeletion in 19q13.2 associated with mental retardation, skeletal malformations, and Diamond-Blackfan anaemia suggests a novel contiguous gene syndrome, **128**
Localisation of the gene causing diaphyseal dysplasia Camurati-Engelmann to chromosome 19q13, **245**
chromosome 1p36, Suggestive evidence for a site specific prostate cancer gene on chromosome 1p36: *letter,* **947**
chromosome 21, Partial tetrasomy 21 in a male infant: *letter,* **e30**
chromosome 2p12-13, Alström syndrome: confirmation of linkage to chromosome 2p12-13 and phenotypic heterogeneity in three affected sibs: *letter,* **e8**
chromosome 3p25, Detailed mapping of a congenital heart disease gene in chromosome 3p25, **581**
chromosome 3q, An analphoid supernumerary marker chromosome derived from chromosome 3 ascertained in a fetus with multiple malformations: *letter,* **807**
chromosome 5, 5p14 deletion associated with microcephaly and seizures, **125**
chromosome 5q, Genetic heterogeneity of Usher syndrome type II: localisation to chromosome 5q, **256**
chromosome 6q22.1→6q23.3, Dysmorphic sibs trisomic for the region 6q22.1→6q23.3: *letter,* **889**
chromosome 7, A novel genetic locus for low renin hypertension: familial hyperaldosteronism type II maps to chromosome 7 (7p22), **831**
chromosome 8, Chromosome 8p alterations in sporadic and *BRCA2* 999del5 linked breast cancer, **342**
chromosome 9, Exclusion of chromosome 9 helps to identify mild variants of acromesomelic dysplasia Maroteaux type, **52**
chromosome marker X22, Detection of fetal cells in transcervical samples using X22 marker: *letter,* **e1**
chromosome regions 2q, 3p, and 10q, A de novo complex chromosomal rearrangement involving chromosomes 2, 3, and 10 associated with microcephaly and early onset spasticity: *letter,* **892**
cilia, A locus for primary ciliary dyskinesia maps to chromosome 19q, **241**
circadian rhythms, Circadian rhythm abnormalities of melatonin in Smith-Magenis syndrome, **428**
cirrhosis, Phosphorylase kinase deficient liver glycogenosis: progression to cirrhosis in infancy associated with *PHKG2* mutations (H144Y and L225R): *letter,* **376**
cleft lip/palate, Partial trisomy 22 in a liveborn resulting from a rearrangement between chromosomes 6 and 22: *letter,* **e22**
clinical facilities, A survey of the current clinical facilities for the management of familial cancer in Europe, **605**
clinical guidelines, Evidence based medicine in practice: lessons from a Scottish clinical genetics project, **684**
clinical history, The Smith-Lemli-Opitz syndrome: *review,* **321**
clinical practice, Genetic registers in clinical practice: a survey of UK clinical geneticists: *letter,* **636**
clinical practice and attitudes, Clinical geneticists' attitudes and practice towards testing for breast cancer susceptibility genes: *letter,* **157**
club foot, Genetics of club foot in Maori and Pacific people, **680**
coarse facies, Hall-Riggs syndrome: a possible second affected family?: *letter,* **886**
cobalamin metabolism, Complementation studies in the *cblA* class of inborn error of cobalamin metabolism: evidence for interallelic complementation and for a new complementation class (*cblH*), **510**
Cockayne syndrome, Cockayne syndrome associated with low CSF 5-hydroxyindole acetic acid levels: *letter,* **553**
Cohen syndrome, Bardet-Biedl and Cohen syndromes: differential diagnostic criteria: *letter,* **e46**
COL10A1, Equal expression of type X collagen mRNA from mutant and wild type *COL10A1* alleles in growth plate cartilage from a patient with metaphyseal chondrodysplasia type Schmid: *letter,* **627**
COL1A2, Glycine to tryptophan substitution in type I collagen in a patient with OI type III: a unique collagen mutation, **371**
***COL2A1* haploinsufficiency mutation,** Clinical variability of Stickler syndrome with a *COL2A1* haploinsufficiency mutation: implications for genetic counselling: *letter,* **318**
COL2A1, Report of five novel and one recurrent *COL2A1* mutations with analysis of genotype-phenotype correlation in patients with a lethal type II collagen disorder, **263**
COL4A5, Somatic mosaicism associated with a mild Alport syndrome phenotype: *letter,* **238**
collagen, Glycine to tryptophan substitution in type I collagen in a patient with OI type III: a unique collagen mutation, **371**
collagen mutations, Clinical phenotypes and molecular characterisation of three patients with Ehlers-Danlos syndrome type VII: *letter,* **e33**
colorectal cancer, Mutational germline analysis of *hMSH2* and *hMLH1* genes in early onset colorectal cancer patients: *letter,* **e7**
The role of hypermethylation of the *hMLH1* promoter region in HNPCC versus MSI+ sporadic colorectal cancers, **588**
complementation class, Complementation studies in the *cblA* class of inborn error of cobalamin metabolism: evidence for interallelic complementation and for a new complementation class (*cblH*), **510**
complex chromosomal rearrangement, A de novo complex chromosomal rearrangement involving chromosomes 2, 3, and 10 associated with microcephaly and early onset spasticity: *letter,* **892**
complex I deficiency, A novel mutation in the thiamine responsive megaloblastic anaemia gene *SLC19A2* in a patient with deficiency of respiratory chain complex I, **669**
complex *CFTR* allele, Fetal bowel hyperechogenicity may indicate mild atypical cystic fibrosis: a case associated with a complex *CFTR* allele: *letter,* **e15**
complex segregation analysis, Genetics of club foot in Maori and Pacific people, **680**
computers, Efficacy of a touchscreen computer based family cancer history questionnaire and subsequent cancer risk assessment, **354**
congenital, The genetics of childhood cataract: *review,* **481**
congenital adrenal hyperplasia, Evidence for digenic inheritance in some cases of Antley-Bixler syndrome?, **26**
congenital disorders of glycosylation, Congenital disorders of glycosylation IIa cause growth retardation, mental retardation, and facial dysmorphism: *letter,* **875**
congenital glaucoma, Molecular characterisation of congenital glaucoma in a consanguineous Canadian community: a step towards preventing glaucoma related blindness, **422**
congenital heart disease, Detailed mapping of a congenital heart disease gene in chromosome 3p25, **581**
connexin 26, A connexin 26 mutation causes a syndrome of sensorineural hearing loss and palmoplantar hyperkeratosis (MIM 148350), **50**
Prevalent connexin 26 gene (*GJB2*) mutations in Japanese, **41**
connexin26, Determination of the frequency of connexin26 mutations in inherited sensorineural deafness and carrier rates in the Tunisian population using DGGE: *letter,* **e39**
connexin26 gene (*GJB2*), A novel C202F mutation in the connexin26 gene (*GJB2*) associated with autosomal dominant isolated hearing loss, **368**
COPS3, Circadian rhythm abnormalities of melatonin in Smith-Magenis syndrome, **428**
Cowden syndrome, Absence of germline mutations in *MINPP1*, a phosphatase encoding gene centromeric of *PTEN*, in patients with Cowden and Bannayan-Riley- Ruvalcaba syndrome without germline *PTEN* mutations: *letter,* **715**
Will the real Cowden syndrome please stand up: revised diagnostic criteria: *commentary,* **828**
craniofacial dysmorphism, Punctate calcification of the epiphyses, visceral malformations, and craniofacial dysmorphism in a female baby: *letter,* **796**
CREB binding protein (CBP/CREBBP), Diagnostic analysis of Rubinstein-Taybi syndrome: five cosmids should be used for microdeletion detection and low number of protein truncating mutations, **168**
cri du chat syndrome, 5p14 deletion associated with microcephaly and seizures, **125**
The first three mosaic cri du chat syndrome patients with two rearranged cell lines: *letter,* **967**
Crigler-Najjar syndrome, Crigler-Najjar syndrome type II resulting from three different mutations in the bilirubin uridine 5'-diphosphate-glucuronosyltransferase (*UGT1A1*) gene: *letter,* **712**
CUL2, Familial clear cell renal cell carcinoma (FCRC): clinical features and mutation analysis of the *VHL*, *MET*, and *CUL2* candidate genes, **348**
Currarino triad, Autosomal dominant sacral agenesis: Currarino syndrome: *review,* **561**
CYP1B1, Molecular characterisation of congenital glaucoma in a consanguineous Canadian community: a step towards preventing glaucoma related blindness, **422**
CYP21 deficiency, Evidence for digenic inheritance in some cases of Antley-Bixler syndrome?, **26**
cystic fibrosis, A novel mutation in the *CFTR* gene correlates with severe clinical phenotype in seven Hispanic patients: *letter,* **215**
Correlation between mutations and age in cystic fibrosis in a French Canadian population: *letter,* **225**
Distribution of *CFTR* gene mutations in cystic fibrosis patients from Estonia: *letter,* **e16**

Fetal bowel hyperechogenicity may indicate mild atypical cystic fibrosis: a case associated with a complex *CFTR* allele: *letter*, **e15**
Genotype-phenotype correlation in three homozygotes and nine compound heterozygotes for the cystic fibrosis mutation 2183AA→G shows a severe phenotype: *letter*, **307**
Many ΔF508 heterozygote neonates with transient hypertrypsinaemia have a second, mild *CFTR* mutation: *letter*, **543**
The relationship between neonatal immunoreactive trypsinogen, ΔF508, and IVS8-5T: *letter*, **629**
cystic fibrosis transmembrane conductance regulator protein (CFTR), The relationship between neonatal immunoreactive trypsinogen, ΔF508, and IVS8-5T: *letter*, **629**
cytosine methylation, Cytosine methylation confers instability on the cardiac troponin T gene in hypertrophic cardiomyopathy: *letter*, **e18**

ΔF508, The relationship between neonatal immunoreactive trypsinogen, ΔF508, and IVS8-5T: *letter*, **629**
de novo translocation, Unexpected high frequency of de novo unbalanced translocations in patients with Wolf-Hirschhorn syndrome (WHS): *letter*, **798**
deafness, An HDR (hypoparathyroidism, deafness, renal dysplasia) syndrome locus maps distal to the DiGeorge syndrome region on 10p13/14, **33**
Characterisation and genetic mapping of a new X linked deafness syndrome, **836**
Tietz syndrome (hypopigmentation/deafness) caused by mutation of *MITF* , **446**
DEFECT 11, The *ALX4* homeobox gene is mutated in patients with ossification defects of the skull (foramina parietalia permagna, OMIM 168500), **916**
7-dehydrocholesterol reductase, The Smith-Lemli-Opitz syndrome: *review*, **321**
deletion, No evidence for imprinting in distal 18q: *letter*, **152**
deletion (2)(p14p15), Deletion (2)(p14p15) in a child with severe neurodevelopmental delay: *letter*, **e21**
deletion 5q, Deletion and duplication of the adenomatous polyposis coli gene resulting from an interchromosomal insertion involving 5(q22q23.3) in the father: *letter*, **141**
deletion mapping, FISH deletion mapping defines a single location for the Y chromosome stature gene, *GCY*, **593**
deletions, Characterisation of six large deletions in *TSC2* identified using long range PCR suggests diverse mechanisms including Alu mediated recombination: *letter*, **877**
denaturing gradient gel electrophoresis, Mutation analysis in glutaric aciduria type I, **177**
dental abnormalities, Two further cases of Sener syndrome: frontonasal dysplasia and dilated Virchow-Robin spaces: *letter*, **466**
Denys-Drash syndrome, Constitutional *WT1* mutations correlate with clinical features in children with progressive nephropathy: *letter*, **698**
***desmin* splice site mutations,** Desmin splice variants causing cardiac and skeletal myopathy, **851**
DGGE, Determination of the frequency of connexin26 mutations in inherited sensorineural deafness and carrier rates in the Tunisian population using DGGE: *letter*, **e39**
DGS/VCFS patients, Investigation of meiotic rearrangements in DGS/VCFS patients with a microdeletion 22q11.2: *letter*, **452**
diaphragmatic hernia, Sex reversal and diaphragmatic hernia in phenotypically female sibs with normal XY chromosomes: *letter*, **315**
differential diagnosis, Bardet-Biedl and Cohen syndromes: differential diagnostic criteria: *letter*, **e46**
DiGeorge syndrome, An HDR (hypoparathyroidism, deafness, renal dysplasia) syndrome locus maps distal to the DiGeorge syndrome region on 10p13/14, **33**
DiGeorge syndrome with discordant phenotype in monozygotic twins: *letter*, **e23**
dilated Virchow-Robin spaces, Two further cases of Sener syndrome: frontonasal dysplasia and dilated Virchow-Robin spaces: *letter*, **466**
dipeptidyl carboxypeptidase 1, Dipeptidyl carboxypeptidase 1 (*DCP1*) and butyrylcholinesterase (*BCHE*) gene interactions with the apolipoprotein E ε4 allele as risk factors in Alzheimer's disease and in Parkinson's disease with coexisting Alzheimer pathology, **766**
direct duplication, Parental origin and mechanisms of formation of cytogenetically recognisable de novo direct and inverted duplications, **281**
direct inheritance, Chromosomal duplication of band 10p14 segregating through four generations: *letter*, **233**
discordant phenotype, DiGeorge syndrome with discordant phenotype in monozygotic twins: *letter*, **e23**
disease, Chromatin modification and disease: *review*, **905**
disease modifiers, Expression of HCM causing mutations: lessons learnt from genotype-phenotype studies of the South African founder *MYH7* A797T mutation: *letter*, **951**
distal arthrogryposis, Progressive neurological deterioration in a child with distal arthrogryposis and whistling face: *letter*, **231**
distal myopathy, Distal spinal muscular atrophy with vocal cord paralysis (dSMA-VII) is not linked to the *MPD2* locus on chromosome 5q31: *letter*, **e14**
distal spinal muscular atrophy, Distal spinal muscular atrophy with vocal cord paralysis (dSMA-VII) is not linked to the *MPD2* locus on chromosome 5q31: *letter*, **e14**
distal trisomy 2p, Distal trisomy 2p and arachnodactyly: *letter*, **974**
distichiasis, Reduction of the genetic interval for lymphoedema-distichiasis to below 2 Mb: *letter*, **725**
DNA binding activity, *ATR-X* mutations cause impaired nuclear location and altered DNA binding properties of the XNP/ATR-X protein, **746**
DNA testing, Clinical geneticists' attitudes and practice towards testing for breast cancer susceptibility genes: *letter*, **157**
Down syndrome, Partial tetrasomy 21 in a male infant: *letter*, **e30**
Duane syndrome, Abnormal sex differentiation and multiple congenital abnormalities in a subject harbouring an apparently balanced (6;8) translocation: *letter*, **972**
duplication, Chromosomal duplication of band 10p14 segregating through four generations: *letter*, **233**
duplication 15q, Duplication of medial 15q confirmed by FISH: *letter*, **e10**
duplication 5q, Deletion and duplication of the adenomatous polyposis coli gene resulting from an interchromosomal insertion involving 5(q22q23.3) in the father: *letter*, **141**
dwarfism, A case of dyschondrosteosis from Roman Britain: *letter*, **e27**
dyschondrosteosis, A case of dyschondrosteosis from Roman Britain: *letter*, **e27**
dyslexia, Two translocations of chromosome 15q associated with dyslexia, **771**
dysmorphic syndrome, New MR/MCA syndrome with distinct facial appearance and general habitus, broad and webbed neck, hypoplastic inverted nipples, epilepsy, and pachygyria of the frontal lobes: *letter*, **460**
dystrophin associated proteins, Disruption of heart sarcoglycan complex and severe cardiomyopathy caused by β sarcoglycan mutations, **102**

ear, Breast hypoplasia and disproportionate short stature in the ear, patella, short stature syndrome: expansion of the phenotype?: *letter*, **719**
early onset breast cancer, A case of inv dup(8p) with early onset breast cancer: *letter*, **70**
Low prevalence of germline *BRCA1* mutations in early onset breast cancer without a family history: *letter*, **792**
ectodermal dysplasia, Pili torti et canaliculi and agenesis of the teeth: report of a new "pure" hair-tooth ectodermal dysplasia in a Norwegian family: *letter*, **721**
ectrodactyly, Delineation of a complex karyotypic rearrangement by microdissection and CGH in a family affected with split foot, **442**
Ehlers-Danlos syndrome type VII (arthrochalasia type), Clinical phenotypes and molecular characterisation of three patients with Ehlers-Danlos syndrome type VII: *letter*, **e33**
elbow dysplasia, Liebenberg syndrome: brachydactyly with joint dysplasia (MIM 186550): a second family: *letter*, **548**
endometriosis, Delineation of a new syndrome: clustering of pyloric stenosis, endometriosis, and breast cancer in two families: *letter*, **794**
epigenotype-phenotype correlations, Epigenotype-phenotype correlations in Beckwith-Wiedemann syndrome, **921**
epilepsy, Interstitial deletion in Xp22.3 is associated with X linked ichthyosis, mental retardation, and epilepsy, **600**
New MR/MCA syndrome with distinct facial appearance and general habitus, broad and webbed neck, hypoplastic inverted nipples, epilepsy, and pachygyria of the frontal lobes: *letter*, **460**
ERCC, Rough skin, brittle hair, and photosensitivity: a mild phenotypic variant of trichothiodystrophy: *letter*, **312**
Europe, A survey of the current clinical facilities for the management of familial cancer in Europe, **605**
evidence based medicine, Evidence based medicine in practice: lessons from a Scottish clinical genetics project, **684**
exon skipping, Clinical phenotypes and molecular characterisation of three patients with Ehlers-Danlos syndrome type VII: *letter*, **e33**
expression study, Desmin splice variants causing cardiac and skeletal myopathy, **851**
***EYA* family genes,** The gene for branchio-oculo-facial syndrome does not colocalise to the *EYA1-4* genes: *letter*, **620**
***EYA1*,** Importance of clinical evaluation and molecular testing in the branchio-oto-renal (BOR) syndrome and overlapping phenotypes: *letter*, **623**
eyaHR, Importance of clinical evaluation and molecular testing in the branchio-oto-renal (BOR) syndrome and overlapping phenotypes: *letter*, **623**

facial clefting, The first description of lethal pterygium syndrome with facial clefting (Bartsocas-Papas syndrome) in 1600: *letter*, **314**
factor V Leiden mutation, Hemiplegic cerebral palsy and the factor V Leiden mutation: *letter*, **787**
familial adenomatous polyposis, Appendiceal carcinoma complicating adenomatous polyposis in a young woman with a de novo constitutional reciprocal translocation t(5;8)(q22;p23.1): *letter*, **71**

familial breast cancer, Why do women attend familial breast cancer clinics?, **197**
familial breast cancer clinics, Why do women attend familial breast cancer clinics?, **197**
familial cancer, A survey of the current clinical facilities for the management of familial cancer in Europe, **605**
familial clear cell renal carcinoma, Familial clear cell renal cell carcinoma (FCRC): clinical features and mutation analysis of the *VHL*, *MET*, and *CUL2* candidate genes, **348**
familial hyperaldosteronism type II, A novel genetic locus for low renin hypertension: familial hyperaldosteronism type II maps to chromosome 7 (7p22), **831**
familial hypercholesterolaemia, Universal primer quantitative fluorescent multiplex (UPQFM) PCR: a method to detect major and minor rearrangements of the low density lipoprotein receptor gene, **272**
familial Wilms tumour, Familial Wilms tumour resulting from *WT1* mutation: intronic polymorphism causing artefactual constitutional homozygosity: *letter*, **377**
family history, A model protocol evaluating the introduction of genetic assessment for women with a family history of breast cancer, **192**
The psychological impact of a cancer family history questionnaire completed in general practice: *letter*, **470**
Feingold syndrome, Interstitial deletion of chromosome 17 (del(17)(q22q23.3 confirms a link with oesophageal atresia: *letter*, **701**
fetal anticonvulsant syndrome, A clinical study of 57 children with fetal anticonvulsant syndromes, **489**
fetal bowel hyperechogenicity, Fetal bowel hyperechogenicity may indicate mild atypical cystic fibrosis: a case associated with a complex *CFTR* allele: *letter*, **e15**
fetal cells, Comorbidity of 5,10-methylenetetrahydrofolate reductase and methionine synthase gene polymorphisms and risk for neural tube defects: *letter*, **949**
Detection of fetal cells in transcervical samples using X22 marker: *letter*, **e1**
fetal valproate syndrome, A clinical study of 57 children with fetal anticonvulsant syndromes, **489**
FGFR3, Achondroplasia with the *FGFR3* 1138g→a (G380R) mutation in two sibs sharing a 4p haplotype derived from their unaffected father: *letter*, **958**
FGFR, Evidence for digenic inheritance in some cases of Antley-Bixler syndrome?, **26**
fibre-FISH, A microdeletion in 19q13.2 associated with mental retardation, skeletal malformations, and Diamond-Blackfan anaemia suggests a novel contiguous gene syndrome, **128**
fibrillin, The molecular genetics of Marfan syndrome and related microfibrillopathies: *review*, **9**
fibroblast growth factor receptor 3, Germline and somatic mosaicism in achondroplasia: *letter*, **956**
fibroblast growth factor receptor 3 (*FGFR3*) gene, Clinical and radiographic features of a family with hypochondroplasia owing to a novel Asn540Scr mutation in the fibroblast growth factor receptor 3 gene: *letter*, **220**
fibrous dysplasia, Familial craniofacial fibrous dysplasia: absence of linkage to *GNAS1* and the gene for cherubism: *letter*, **e37**
Finland, Epidemiology of neurofibromatosis type 1 (NF1) in northern Finland: *letter*, **632**
FISH, A supernumerary marker chromosome originating from two different regions of chromosome 18, **121**
Duplication of medial 15q confirmed by FISH: *letter*, **e10**
FISH deletion mapping defines a single location for the Y chromosome stature gene, *GCY* , **593**
NF2 gene deletion in a family with a mild phenotype: *letter*, **75**
Molecular cytogenetic characterisation of partial trisomy 9q in a case with pyloric stenosis and a review, **529**
Partial trisomy 22 in a liveborn resulting from a rearrangement between chromosomes 6 and 22: *letter*, **e22**
Perfect endings: a review of subtelomeric probes and their use in clinical diagnosis: *review*, **401**
fluorescence in situ hybridisation, 5p14 deletion associated with microcephaly and seizures, **125**
Identification of supernumerary marker chromosomes derived from chromosomes 5, 6, 19, and 20 using FISH, **114**
Prenatal detection of trisomy for the entire long arm of chromosome 7: *letter*, **551**
FMR2, *FMR3* is a novel gene associated with *FRAXE* CpG island and transcriptionally silent in *FRAXE* full mutations, **782**
FMR3, *FMR3* is a novel gene associated with *FRAXE* CpG island and transcriptionally silent in *FRAXE* full mutations, **782**
FMRP, Increase of FMRP expression, raised levels of *FMR1* mRNA, and clonal selection in proliferating cells with unmethylated fragile X repeat expansions: a clue to the sex bias in the transmission of full mutations?, **842**
founder mutation, Recurrent germline mutation in *MSH2* arises frequently de novo, **646**
FPP, The *ALX4* homeobox gene is mutated in patients with ossification defects of the skull (foramina parietalia permagna, OMIM 168500), **916**
fragile X syndrome, Absence of fragile X syndrome in Nova Scotia: *letter*, **77**
Increase of FMRP expression, raised levels of *FMR1* mRNA, and clonal selection in proliferating cells with unmethylated fragile X repeat expansions: a clue to the sex bias in the transmission of full mutations?, **842**
Twin sisters, monozygotic with the fragile X mutation, but with a different phenotype, **603**
Frasier syndrome, Constitutional *WT1* mutations correlate with clinical features in children with progressive nephropathy: *letter*, **698**
FRAX syndromes, FRAXA and FRAXE: the results of a five year survey, **415**
FRAXE, *FMR3* is a novel gene associated with *FRAXE* CpG island and transcriptionally silent in *FRAXE* full mutations, **782**
***FRDA* frameshift mutations,** Identification of three novel frameshift mutations in patients with Friedreich's ataxia: *letter*, **e38**
***FRDA* gene,** Friedreich ataxia: an overview: *review*, **1**
Freeman-Sheldon syndrome, Progressive neurological deterioration in a child with distal arthrogryposis and whistling face: *letter*, **231**
frequencies, Prevalence of mitochondrial gene mutations among hearing impaired patients, **38**
frequency, Disease associated balanced chromosome rearrangements: a resource for large scale genotype-phenotype delineation in man, **858**
Friedreich ataxia, Friedreich ataxia: an overview: *review*, **1**
Friedreich's ataxia, Identification of three novel frameshift mutations in patients with Friedreich's ataxia: *letter*, **e38**
frontonasal dysplasia, Two further cases of Sener syndrome: frontonasal dysplasia and dilated Virchow-Robin spaces: *letter*, **466**

GABRR1, Mutation analysis of *GABRR1* and *GABRR2* in autosomal recessive retinitis pigmentosa (*RP25*): *letter*, **e5**
GABRR2, Mutation analysis of *GABRR1* and *GABRR2* in autosomal recessive retinitis pigmentosa (*RP25*): *letter*, **e5**
G→C splice acceptor site mutation, Incidence and molecular mechanism of aberrant splicing owing to a G→C splice acceptor site mutation causing Smith-Lemli-Opitz syndrome: *letter*, **387**
Gaucher disease, Cardiovascular fibrosis, hydrocephalus, ophthalmoplegia, and visceral involvement in an American child with Gaucher disease: *letter*, **e40**
GCY, FISH deletion mapping defines a single location for the Y chromosome stature gene, *GCY* , **593**
gene, Molecular characterisation of congenital glaucoma in a consanguineous Canadian community: a step towards preventing glaucoma related blindness, **422**
gene deletion, *NF2* gene deletion in a family with a mild phenotype: *letter*, **75**
genetic assessment, A model protocol evaluating the introduction of genetic assessment for women with a family history of breast cancer, **192**
genetic counselling, An evaluation of needs of female *BRCA1* and *BRCA2* carriers undergoing genetic counselling, **866**
Clinical variability of Stickler syndrome with a *COL2A1* haploinsufficiency mutation: implications for genetic counselling: *letter*, **318**
Efficacy of a touchscreen computer based family cancer history questionnaire and subsequent cancer risk assessment, **354**
Evaluation of a counselling protocol for predictive genetic testing for hereditary non-polyposis colorectal cancer, **108**
Genetics and deafness: what do families want?: *letter*, **e26**
Molecular characterisation of congenital glaucoma in a consanguineous Canadian community: a step towards preventing glaucoma related blindness, **422**
Mosaicism in Alport syndrome and genetic counselling: *letter*, **717**
Shared decision making and non-directiveness in genetic counselling, **135**
genetic heterogeneity, Benign familial infantile convulsions: report of a UK family and confirmation of genetic heterogeneity: *letter*, **e31**
genetic homogeneity, The continuing failure to recognise Alström syndrome and further evidence of genetic homogeneity: *letter*, **219**
genetic linkage analysis, A cross section of autosomal recessive limb-girdle muscular dystrophies in 38 families, **361**
genetic predisposition, No evidence of germline *PTEN* mutations in familial prostate cancer: *letter*, **210**
genetic registers, Genetic registers in clinical practice: a survey of UK clinical geneticists: *letter*, **636**
genetic risk, Guidelines for a genetic risk based approach to advising women with a family history of breast cancer, **203**
genetic susceptibility, Genetic susceptibility to age related macular degeneration: *review*, **83**
genetics, First description of germline mosaicism in familial hypertrophic cardiomyopathy, **132**
Identification of cathepsin C mutations in ethnically diverse Papillon-Lefèvre syndrome patients, **927**
Novel mutation in the *MYOC* gene in primary open angle glaucoma patients: *letter*, **301**
The genetics of childhood cataract: *review*, **481**
The heritability of high myopia: a reanalysis of Goldschmidt's data: *letter*, **227**

genetics clinic, Why patients do not attend for their appointments at a genetics clinic: *letter,* **810**
genital anomalies, Genitopatellar syndrome: a new condition comprising absent patellae, scrotal hypoplasia, renal anomalies, facial dysmorphism, and mental retardation, **520**
genotype-phenotype correlation, Desmin splice variants causing cardiac and skeletal myopathy, **851**
Five cases of isolated glycerol kinase deficiency, including two families: failure to find genotype:phenotype correlation, **434**
Genotype-phenotype relationship of Niemann-Pick disease type C: a possible correlation between clinical onset and levels of NPC1 protein in isolated skin fibroblasts: *letter,* **707**
Expression of HCM causing mutations: lessons learnt from genotype-phenotype studies of the South African founder *MYH7* A797T mutation: *letter,* **951**
Genotype-phenotype correlations in tuberous sclerosis: *letter,* **e3**
germline *p16^{INK4a}* alterations, Absence of germline *p16^{INK4a}* alterations in *p53* wild type Li-Fraumeni syndrome families: *letter,* **e13**
germline mosaic, Germline and somatic mosaicism in achondroplasia: *letter,* **956**
germline mosaicism, First description of germline mosaicism in familial hypertrophic cardiomyopathy, **132**
Germline mosaicism for a *GNAS1* mutation and Albright hereditary osteodystrophy: *letter,* **e35**
germline mutation, Cryptic von Hippel-Lindau disease: germline mutations in patients with haemangioblastoma only, **939**
germline mutations, *BRCA2* germline mutations among early onset breast cancer patients unselected for family history of the disease: *letter,* **e17**
Novel germline mutations in the *PTEN* tumour suppressor gene found in women with multiple cancers, **336**
GFRα-3, Characterisation of the human *GFRα-3* locus and investigation of the gene in Hirschsprung disease, **674**
GJB2, Prevalent connexin 26 gene (*GJB2*) mutations in Japanese, **41**
glaucoma, Novel mutation in the *MYOC* gene in primary open angle glaucoma patients: *letter,* **301**
glutaric aciduria type I, Mutation analysis in glutaric aciduria type I, **177**
glutaryl-CoA dehydrogenase, Mutation analysis in glutaric aciduria type I, **177**
glycerol kinase deficiency, Five cases of isolated glycerol kinase deficiency, including two families: failure to find genotype:phenotype correlation, **434**
glycogen storage disease, Phosphorylase kinase deficient liver glycogenosis: progression to cirrhosis in infancy associated with *PHKG2* mutations (H144Y and L225R): *letter,* **376**
glycosylation IIa, Congenital disorders of glycosylation IIa cause growth retardation, mental retardation, and facial dysmorphism: *letter,* **875**
GNAS1, Familial craniofacial fibrous dysplasia: absence of linkage to *GNAS1* and the gene for cherubism: *letter,* **e37**
Germline mosaicism for a *GNAS1* mutation and Albright hereditary osteodystrophy: *letter,* **e35**
gonadal dysgenesis, Abnormal sex differentiation and multiple congenital abnormalities in a subject harbouring an apparently balanced (6;8) translocation: *letter,* **972**
guidelines, Guidelines for a genetic risk based approach to advising women with a family history of breast cancer, **203**

haemangioblastoma, Cryptic von Hippel-Lindau disease: germline mutations in patients with haemangioblastoma only, **939**
Risk of multisystem disease in isolated ocular angioma (haemangioblastoma): *letter,* **62**
Haim-Munk syndrome, Haim-Munk syndrome and Papillon-Lefèvre syndrome are allelic mutations in cathepsin C, **88**
Of palms, soles, and gums: *commentary,* **81**
hair, Can hair be used to screen for breast cancer?: *letter,* **297**
half cryptic translocation, Monosomy and trisomy 1q44-qter in two sisters originating from a half cryptic 1q;15p translocation: *letter,* **612**
Hall-Riggs syndrome, Hall-Riggs syndrome: a possible second affected family?: *letter,* **886**
haplotypes, Alkaptonuria in Italy: polymorphic haplotype background, mutational profile, and description of four novel mutations in the homogentisate 1,2-dioxygenase gene: *letter,* **309**
hearing impairment, A novel mutation in the mitochondrial tRNA$^{Ser(UCN)}$ gene in a family with non-syndromic sensorineural hearing impairment, **692**
Genetics and deafness: what do families want?: *letter,* **e26**
Prevalence of mitochondrial gene mutations among hearing impaired patients, **38**
hearing loss, Genetic heterogeneity of Usher syndrome type II: localisation to chromosome 5q, **256**
heparan-N-sulphatase, Mutational analysis of Sanfilippo syndrome type A (MPS IIIA): identification of 13 novel mutations: *letter,* **704**
hereditary breast cancer, Attitudes to genetic testing for breast cancer susceptibility in women at increased risk of developing hereditary breast cancer: *letter,* **472**
hereditary diseases, Channelopathies: ion channel defects linked to heritable clinical disorders: *review,* **729**
hereditary nephritis, Somatic mosaicism associated with a mild Alport syndrome phenotype: *letter,* **238**
hereditary neuropathy with liability to pressure palsies, Hereditary neuropathy with liability to pressure palsies: two cases with a reciprocal translocation t(16;17)(q12;p11.2) interrupting the *PMP22* gene: *letter,* **396**
hereditary non-polyposis colorectal cancer, Evaluation of a counselling protocol for predictive genetic testing for hereditary non-polyposis colorectal cancer, **108**
Incidence of germline *hMLH1* and *hMSH2* mutations (HNPCC patients) among newly diagnosed colorectal cancers in a Slovenian population: *letter,* **533**
Mutation analysis of *SMAD2*, *SMAD3*, and *SMAD4* genes in hereditary non-polyposis colorectal cancer: *letter,* **298**
Sensitivity and specificity of clinical criteria for hereditary non-polyposis colorectal cancer associated mutations in *MSH2* and *MLH1* , **641**
Two common forms of the human *MLH1* gene may be associated with functional differences, **776**
hereditary pancreatitis, Mutational analysis of the human pancreatic secretory trypsin inhibitor (*PSTI*) gene in hereditary and sporadic chronic pancreatitis: *letter,* **67**
hereditary spastic paraparesis, Mutation analysis of the spastin gene (*SPG4*) in patients with hereditary spastic paraparesis, **759**
heritability, The heritability of high myopia: a reanalysis of Goldschmidt's data: *letter,* **227**
HERV, Divergent outcomes of intrachromosomal recombination on the human Y chromosome: male infertility and recurrent polymorphism, **752**
***HGO* gene,** Novel mutations in the homogentisate-1,2-dioxygenase gene identified in Slovak patients with alkaptonuria: *letter,* **539**
***HGO* mutations,** Novel mutations in the homogentisate-1,2-dioxygenase gene identified in Slovak patients with alkaptonuria: *letter,* **539**
HI, An unbalanced submicroscopic translocation t(8;16)(q24.3;p13.3)pat associated with tuberous sclerosis complex, adult polycystic kidney disease, and hypomelanosis of Ito, **287**
Hirschsprung disease, Characterisation of the human *GFRα-3* locus and investigation of the gene in Hirschsprung disease, **674**
***hMLH1* mutation,** Incidence of germline *hMLH1* and *hMSH2* mutations (HNPCC patients) among newly diagnosed colorectal cancers in a Slovenian population: *letter,* **533**
***hMLH1* promoter region,** The role of hypermethylation of the *hMLH1* promoter region in HNPCC versus MSI+ sporadic colorectal cancers, **588**
hMLH1, Mutational germline analysis of *hMSH2* and *hMLH1* genes in early onset colorectal cancer patients: *letter,* **e7**
***hMSH2* mutation,** Incidence of germline *hMLH1* and *hMSH2* mutations (HNPCC patients) among newly diagnosed colorectal cancers in a Slovenian population: *letter,* **533**
hMSH2, Mutational germline analysis of *hMSH2* and *hMLH1* genes in early onset colorectal cancer patients: *letter,* **e7**
HNPCC, Evaluation of a counselling protocol for predictive genetic testing for hereditary non-polyposis colorectal cancer, **108**
The role of hypermethylation of the *hMLH1* promoter region in HNPCC versus MSI+ sporadic colorectal cancers, **588**
Holt-Oram syndrome, The mutation spectrum in Holt-Oram syndrome: *letter,* **785**
homogentisate 1,2 dioxygenase, Alkaptonuria in Italy: polymorphic haplotype background, mutational profile, and description of four novel mutations in the homogentisate 1,2-dioxygenase gene: *letter,* **309**
homozygosity by descent, Confirmation of the assignment of the Sanjad-Sakati (congenital hypoparathyroidism) syndrome (OMIM 241410) locus to chromosome 1q42-43: *letter,* **63**
homozygosity mapping, Exclusion of chromosome 9 helps to identify mild variants of acromesomelic dysplasia Maroteaux type, **52**
HOX11L1 gene, Evaluation of the *HOX11L1* gene as a candidate for congenital disorders of intestinal innervation: *letter,* **e9**
human chromosome 6, Abnormal sex differentiation and multiple congenital abnormalities in a subject harbouring an apparently balanced (6;8) translocation: *letter,* **972**
human *MLH1* gene, Two common forms of the human *MLH1* gene may be associated with functional differences, **776**
human pancreatic secretory trypsin inhibitor, Mutational analysis of the human pancreatic secretory trypsin inhibitor (*PSTI*) gene in hereditary and sporadic chronic pancreatitis: *letter,* **67**
Huntington's desease Ten years of presymptomatic testing for Huntington's disease; the experience of the UK Huntington's Disease Prediction Consortium, **567**
hydrocephalus, Cardiovascular fibrosis, hydrocephalus, ophthalmoplegia, and visceral involvement in an American child with Gaucher disease: *letter,* **e40**
5-hydroxyindole acetic acid, Cockayne syndrome associated with low CSF 5-hydroxyindole acetic acid levels: *letter,* **553**
5-hydroxytryptophan, Cockayne syndrome associated with low CSF 5-hydroxyindole acetic acid levels: *letter,* **553**

hyperbilirubinaemia, Crigler-Najjar syndrome type II resulting from three different mutations in the bilirubin uridine 5'-diphosphate-glucuronosyltransferase (*UGT1A1*) gene: *letter*, **712**
hypercholesterolaemia, Predominance of a 6 bp deletion in exon 2 of the LDL receptor gene in Africans with familial hypercholesterolaemia, **514**
hyperkeratosis, Identification of cathepsin C mutations in ethnically diverse Papillon-Lefèvre syndrome patients, **927**
hypermethylation, The role of hypermethylation of the *hMLH1* promoter region in HNPCC versus MSI+ sporadic colorectal cancers, **588**
hypertension, A novel genetic locus for low renin hypertension: familial hyperaldosteronism type II maps to chromosome 7 (7p22), **831**
hypertrophic cardiomyopathy, Cytosine methylation confers instability on the cardiac troponin T gene in hypertrophic cardiomyopathy: *letter*, **e18**
Expression of HCM causing mutations: lessons learnt from genotype-phenotype studies of the South African founder *MYH7* A797T mutation: *letter*, **951**
First description of germline mosaicism in familial hypertrophic cardiomyopathy, **132**
hypochondroplasia, Clinical and radiographic features of a family with hypochondroplasia owing to a novel Asn540Ser mutation in the fibroblast growth factor receptor 3 gene: *letter*, **220**
hypomelanosis of Ito, Mosaicism for a dup(12)(q22q13) in a patient with hypomelanosis of Ito and asymmetry: *letter*, **804**
hypoparathyroidism, An HDR (hypoparathyroidism, deafness, renal dysplasia) syndrome locus maps distal to the DiGeorge syndrome region on 10p13/14, **33**
Confirmation of the assignment of the Sanjad-Sakati (congenital hypoparathyroidism) syndrome (OMIM 241410) locus to chromosome 1q42-43: *letter*, **63**

I allele, The *ACE* gene and Alzheimer's disease susceptibility, **695**
idiopathic multicentric osteolysis, Idiopathic multicentric osteolysis presents early and is not linked to chromosome 18q21.1: *letter*, **e34**
***IGFBP1*,** Biallelic expression of *IGFBP1* and *IGFBP3*, two candidate genes for the Silver-Russell syndrome: *letter*, **65**
***IGFBP3*,** Biallelic expression of *IGFBP1* and *IGFBP3*, two candidate genes for the Silver-Russell syndrome: *letter*, **65**
***IL-4Rα* gene,** No linkage or association of the *IL-4Rα* gene Q576R mutation with atopic asthma in Italian families: *letter*, **382**
immunoreactive trypsinogen, Many ΔF508 heterozygote neonates with transient hypertrypsinaemia have a second, mild *CFTR* mutation: *letter*, **543**
immunoreactive trypsinogen (IRT), The relationship between neonatal immunoreactive trypsinogen, ΔF508, and IVS8-5T: *letter*, **629**
imperforate anus, Partial trisomy 22 in a liveborn resulting from a rearrangement between chromosomes 6 and 22: *letter*, **e22**
imprinting, Biallelic expression of *IGFBP1* and *IGFBP3*, two candidate genes for the Silver-Russell syndrome: *letter*, **65**
Epigenotype-phenotype correlations in Beckwith-Wiedemann syndrome, **921**
Maternal uniparental heterodisomy of chromosome 14: chromosomal mechanism and clinical follow up, **525**
No evidence for imprinting in distal 18q: *letter*, **152**
in situ hybridisation, Mutation screening in Rett syndrome patients, **250**
incidence, Epidemiology of neurofibromatosis type 1 (NF1) in northern Finland: *letter*, **632**
FRAXA and FRAXE: the results of a five year survey, **415**
infantile autism, Spectrum of mutations in the *MECP2* gene in patients with infantile autism and Rett syndrome: *letter*, **e41**
infertility, Abnormal sex differentiation and multiple congenital abnormalities in a subject harbouring an apparently balanced (6;8) translocation: *letter*, **972**
Divergent outcomes of intrachromosomal recombination on the human Y chromosome: male infertility and recurrent polymorphism, **752**
inguinal hernia, Two further cases of Sener syndrome: frontonasal dysplasia and dilated Virchow-Robin spaces: *letter*, **466**
inhibin, Autosomal translocation associated with premature ovarian failure: *letter*, **e2**
insertion, Deletion and duplication of the adenomatous polyposis coli gene resulting from an interchromosomal insertion involving 5(q22q23.3) in the father: *letter*, **141**
intermediate alleles, FRAXA and FRAXE: the results of a five year survey, **415**
interphase FISH, Rapid detection of microdeletions using fluorescence in situ hybridisation (FISH) on buccal smears: *letter*, **e4**
interstitial deletion, 5p14 deletion associated with microcephaly and seizures, **125**
Interstitial deletion of chromosome 17 (del(17)(q22q23.3)) confirms a link with oesophageal atresia: *letter*, **701**
interstitial deletion 16q, Clinical and molecular cytogenetic studies of a large de novo interstitial deletion 16q11.2-16q21 including the putative transcription factor gene *SALL1* : *letter*, **389**
intestinal innervation, Evaluation of the *HOX11L1* gene as a candidate for congenital disorders of intestinal innervation: *letter*, **e9**
intestinal neuronal dysplasia type B (IND B), Tandem duplication within the neurofibromatosis type 1 gene (*NF1*) and reciprocal t(15;16)(q26.3;q12.1) translocation in familial association of NF1 with intestinal neuronal dysplasia type B (IND B): *letter*, **146**
intrauterine growth retardation, Confirmation of the assignment of the Sanjad-Sakati (congenital hypoparathyroidism) syndrome (OMIM 241410) locus to chromosome 1q42-43: *letter*, **63**
inv dup(8p), A case of inv dup(8p) with early onset breast cancer: *letter*, **70**
inversions, Disease associated balanced chromosome rearrangements: a resource for large scale genotype-phenotype delineation in man, **858**
inverted duplication, Parental origin and mechanisms of formation of cytogenetically recognisable de novo direct and inverted duplications, **281**
inverted single band triplication, Tandem triplication of chromosome 13q14 with inverted interstitial segment in a 4 year old girl: *letter*, **964**
ion channel genetics, Channelopathies: ion channel defects linked to heritable clinical disorders: *review*, **729**
ion channel physiopathology, Channelopathies: ion channel defects linked to heritable clinical disorders: *review*, **729**
isochromosome, Pure trisomy 20p resulting from isochromosome formation and whole arm translocation: *letter*, **454**
Italy, Alkaptonuria in Italy: polymorphic haplotype background, mutational profile, and description of four novel mutations in the homogentisate 1,2-dioxygenase gene: *letter*, **309**

***JAGGED*1,** *JAGGED*1 expression in human embryos: correlation with the Alagille syndrome phenotype, **658**
Japanese, Prevalent connexin 26 gene (*GJB2*) mutations in Japanese, **41**

Kartagener syndrome, A locus for primary ciliary dyskinesia maps to chromosome 19q, **241**
Kenny-Caffey syndrome, Hypoparathyroidism, retarded growth and development, and dysmorphism or Sanjad-Sakati syndrome: an Arab disease reminiscent of Kenny-Caffey syndrome: *letter*, **145**

Langer mesomelic dysplasia, Homozygous deletion of *SHOX* in a mentally retarded male with Langer mesomelic dysplasia: *letter*, **959**
***LDLR*,** Universal primer quantitative fluorescent multiplex (UPQFM) PCR: a method to detect major and minor rearrangements of the low density lipoprotein receptor gene, **272**
learning difficulty, Renal angiomyolipomata and learning difficulty in tuberous sclerosis complex: *letter*, **156**
learning disorder, Characterisation and genetic mapping of a new X linked deafness syndrome, **836**
lethal pterygium syndrome, The first description of lethal pterygium syndrome with facial clefting (Bartsocas-Papas syndrome) in 1600: *letter*, **314**
Li-Fraumeni syndrome, Absence of germline $p16^{INK4a}$ alterations in *p53* wild type Li-Fraumeni syndrome families: *letter*, **e13**
limb-girdle muscular dystrophy, A cross section of autosomal recessive limb-girdle muscular dystrophies in 38 families, **361**
Disruption of heart sarcoglycan complex and severe cardiomyopathy caused by β sarcoglycan mutations, **102**
linkage, A locus for primary ciliary dyskinesia maps to chromosome 19q, **241**
Localisation of a gene for prepubertal periodontitis to chromosome 11q14 and identification of a cathepsin C gene mutation, **95**
linkage analysis, Importance of the autosomal recessive retinitis pigmentosa locus on 1q31-q32.1 (RP12) and mutation analysis of the candidate gene *RGS16* (*RGS-r*): *letter*, **384**
No evidence of germline *PTEN* mutations in familial prostate cancer: *letter*, **210**
LOH, Chromosome 8p alterations in sporadic and *BRCA2* 999del5 linked breast cancer, **342**
long range PCR, Characterisation of six large deletions in *TSC2* identified using long range PCR suggests diverse mechanisms including Alu mediated recombination: *letter*, **877**
low density lipoprotein receptor, Predominance of a 6 bp deletion in exon 2 of the LDL receptor gene in Africans with familial hypercholesterolaemia, **514**
low density lipoprotein receptor gene (*LDLR*), I705 variant in the low density lipoprotein receptor gene has no effect on plasma cholesterol levels: *letter*, **713**
low penetrance, A constitutional homozygous mutation in the *RB1* gene in a patient with unilateral retinoblastoma: *letter*, **615**
low penetrance gene, Detecting low penetrance genes in cancer: the way ahead: *review*, **161**
lymphoedema, Reduction of the genetic interval for lymphoedema-distichiasis to below 2 Mb: *letter*, **725**
Lyonisation, Twin sisters, monozygotic with the fragile X mutation, but with a different phenotype, **603**

macrocephaly, Growth in North American white children with neurofibromatosis 1 (NF1), **933**
Madelung deformity, A case of dyschondrosteosis from Roman Britain: *letter,* **e27**
major rearrangements, Universal primer quantitative fluorescent multiplex (UPQFM) PCR: a method to detect major and minor rearrangements of the low density lipoprotein receptor gene, **272**
male breast cancer, The CAG repeat within the androgen receptor gene in male breast cancer patients: *letter,* **139**
management, The Smith-Lemli-Opitz syndrome: *review,* **321**
Marfan syndrome, The molecular genetics of Marfan syndrome and related microfibrillopathies: *review,* **9**
marfanoid, Distal trisomy 2p and arachnodactyly: *letter,* **974**
maternal paracentric inversion, A child with bisatellited, dicentric chromosome 15 arising from a maternal paracentric inversion of chromosome 15q: *letter,* **e11**
maternal uniparental disomy 7, Silver-Russell syndrome and ring chromosome 7: *letter,* **380**
maternal UPD, Maternal uniparental heterodisomy of chromosome 14: chromosomal mechanism and clinical follow up, **525**
MCP, Maternal uniparental heterodisomy of chromosome 14: chromosomal mechanism and clinical follow up, **525**
MECP2, Spectrum of mutations in the *MECP2* gene in patients with infantile autism and Rett syndrome: *letter,* **e41**
***MECP2* gene,** Mutation analysis of the methyl-CpG binding protein 2 gene (*MECP2*) in patients with Rett syndrome: *letter,* **608**
Mutations in the *MECP2* gene in a cohort of girls with Rett syndrome: *letter,* **610**
medical informatics, Efficacy of a touchscreen computer based family cancer history questionnaire and subsequent cancer risk assessment, **354**
medulloblastoma, Gene amplification in PNETs/medulloblastomas: mapping of a novel amplified gene within the *MYCN* amplicon, **501**
meiotic rearrangements, Investigation of meiotic rearrangements in DGS/VCFS patients with a microdeletion 22q11.2: *letter,* **452**
melanoma, Identification of *PTEN* mutations in metastatic melanoma specimens, **653**
melatonin, Circadian rhythm abnormalities of melatonin in Smith-Magenis syndrome, **428**
meningioma, Neurofibromatosis type 2: *review,* **897**
mental retardation, Characterisation and genetic mapping of a new X linked deafness syndrome, **836**
Genitopatellar syndrome: a new condition comprising absent patellae, scrotal hypoplasia, renal anomalies, facial dysmorphism, and mental retardation, **520**
Hall-Riggs syndrome: a possible second affected family?: *letter,* **886**
Homozygous deletion of *SHOX* in a mentally retarded male with Langer mesomelic dysplasia: *letter,* **959**
FMR3 is a novel gene associated with *FRAXE* CpG island and transcriptionally silent in *FRAXE* full mutations, **782**
Progressive neurological deterioration in a child with distal arthrogryposis and whistling face: *letter,* **231**
Twin sisters, monozygotic with the fragile X mutation, but with a different phenotype, **603**
metacarpophalangeal profile, Clinical and radiological assessment of a family with mild brachydactyly type A1: the usefulness of metacarpophalangeal profiles, **292**
metaphyseal chondrodysplasia type Schmid, Equal expression of type X collagen mRNA from mutant and wild type *COL10A1* alleles in growth plate cartilage from a patient with metaphyseal chondrodysplasia type Schmid: *letter,* **627**
methionine synthase, Comorbidity of 5,10-methylenetetrahydrofolate reductase and methionine synthase gene polymorphisms and risk for neural tube defects: *letter,* **949**
methylation specific PCR assay, Diagnosis of maternal uniparental disomy of chromosome 7 with a methylation specific PCR assay: *letter,* **e19**
5,10-methylenetetrahydrofolate reductase, Comorbidity of 5,10-methylenetetrahydrofolate reductase and methionine synthase gene polymorphisms and risk for neural tube defects: *letter,* **949**
MET, Familial clear cell renal cell carcinoma (FCRC): clinical features and mutation analysis of the *VHL, MET,* and *CUL2* candidate genes, **348**
M-FISH, An analphoid supernumerary marker chromosome derived from chromosome 3 ascertained in a fetus with multiple malformations: *letter,* **807**
microcephaly, A de novo complex chromosomal rearrangement involving chromosomes 2, 3, and 10 associated with microcephaly and early onset spasticity: *letter,* **892**
microdeletion, Diagnostic analysis of Rubinstein-Taybi syndrome: five cosmids should be used for microdeletion detection and low number of protein truncating mutations, **168**
microdeletion 22q11.2, Investigation of meiotic rearrangements in DGS/VCFS patients with a microdeletion 22q11.2: *letter,* **452**
microdeletion syndrome, A microdeletion in 19q13.2 associated with mental retardation, skeletal malformations, and Diamond-Blackfan anaemia suggests a novel contiguous gene syndrome, **128**
microdeletion syndromes, Rapid detection of microdeletions using fluorescence in situ hybridisation (FISH) on buccal smears: *letter,* **e4**
microdissection, A supernumerary marker chromosome originating from two different regions of chromosome 18, **121**
Delineation of a complex karyotypic rearrangement by microdissection and CGH in a family affected with split foot, **442**
microfibrillopathies, The molecular genetics of Marfan syndrome and related microfibrillopathies: *review,* **9**
microsatellite instability, Incidence of germline *hMLH1* and *hMSH2* mutations (HNPCC patients) among newly diagnosed colorectal cancers in a Slovenian population: *letter,* **533**
Low frequency of microsatellite instability in *BRCA1* mutated breast tumours: *letter,* **e32**
microsatellite repeat, Identification of novel alleles at a polymorphic microsatellite repeat region in the human *NRAMP1* gene promoter: analysis of allele frequencies in primary biliary cirrhosis: *letter,* **150**
microtia, Breast hypoplasia and disproportionate short stature in the ear, patella, short stature syndrome: expansion of the phenotype?: *letter,* **719**
MINPP1, Absence of germline mutations in *MINPP1,* a phosphatase encoding gene centromeric of *PTEN,* in patients with Cowden and Bannayan-Riley-Ruvalcaba syndrome without germline *PTEN* mutations: *letter,* **715**
missense mutation, Two novel germline mutations of the retinoblastoma gene (*RB1*) that show incomplete penetrance, one splice site and one missense: *letter,* **e6**
MITF, Tietz syndrome (hypopigmentation/deafness) caused by mutation of *MITF* , **446**
mitochondria, Prevalence of mitochondrial gene mutations among hearing impaired patients, **38**
mitochondrial DNA, Further evidence for genetic heterogeneity of autosomal dominant disorders with accumulation of multiple deletions of mitochondrial DNA: *letter,* **547**
mitochondrial genome, The mitochondrial genome in Wolfram syndrome: *letter,* **463**
MLH1, Sensitivity and specificity of clinical criteria for hereditary non-polyposis colorectal cancer associated mutations in *MSH2* and *MLH1* , **641**
modifier genes, Tandem duplication within the neurofibromatosis type 1 gene (*NF1*) and reciprocal t(15;16)(q26.3;q12.1) translocation in familial association of NF1 with intestinal neuronal dysplasia type B (IND B): *letter,* **146**
monosomy 1q, Monosomy and trisomy 1q44-qter in two sisters originating from a half cryptic 1q;15p translocation: *letter,* **612**
monozygotic twins, DiGeorge syndrome with discordant phenotype in monozygotic twins: *letter,* **e23**
Twin sisters, monozygotic with the fragile X mutation, but with a different phenotype, **603**
mosaic, Neurofibromatosis type 2: *review,* **897**
mosaicism, A constitutional homozygous mutation in the *RB1* gene in a patient with unilateral retinoblastoma: *letter,* **615**
Mosaicism in Alport syndrome and genetic counselling: *letter,* **717**
Skin pigmentary anomalies in a mosaic form of partial tetrasomy 3q: *letter,* **723**
MPS II, MPS II in females: molecular basis of two different cases: *letter,* **e29**
MRI, Use of MRI and audiological tests in presymptomatic diagnosis of type 2 neurofibromatosis (NF2): *letter,* **944**
MR/MCA syndrome, New MR/MCA syndrome with distinct facial appearance and general habitus, broad and webbed neck, hypoplastic inverted nipples, epilepsy, and pachygyria of the frontal lobes: *letter,* **460**
MSH2, Sensitivity and specificity of clinical criteria for hereditary non-polyposis colorectal cancer associated mutations in *MSH2* and *MLH1,* **641**
Recurrent germline mutation in *MSH2* arises frequently de novo, **646**
mtDNA mutation, A novel mutation in the mitochondrial $tRNA^{Ser(UCN)}$ gene in a family with non-syndromic sensorineural hearing impairment, **692**
mucopolysaccharidosis IIIA, Mutational analysis of Sanfilippo syndrome type A (MPS IIIA): identification of 13 novel mutations: *letter,* **704**
multicolour chromosomal bar codes, Mosaicism for a dup(12)(q22q13) in a patient with hypomelanosis of Ito and asymmetry: *letter,* **804**
multiple cancers, Novel germline mutations in the *PTEN* tumour suppressor gene found in women with multiple cancers, **336**
multiple endocrine neoplasia type 2, Multiple endocrine neoplasia type 2 and *PET*: from neoplasia to neurogenesis: *review,* **817**
multi-telomere FISH, Cryptic subtelomeric translocations in the 22q13 deletion syndrome, **58**
mutation, Determination of the frequency of connexin26 mutations in inherited sensorineural deafness and carrier rates in the Tunisian population using DGGE: *letter,* **e39**
I705 variant in the low density lipoprotein receptor gene has no effect on plasma cholesterol levels: *letter,* **713**
ATR-X mutations cause impaired nuclear location and altered DNA binding properties of the XNP/ATR-X protein, **746**

Mutation analysis in glutaric aciduria type I, 177
Mutation analysis of the spastin gene (*SPG4*) in patients with hereditary spastic paraparesis, 759
Novel mutation in the *MYOC* gene in primary open angle glaucoma patients: *letter*, 301
Predominance of a 6 bp deletion in exon 2 of the LDL receptor gene in Africans with familial hypercholesterolaemia, 514
Three novel *SALL1* mutations extend the mutational spectrum in Townes-Brocks syndrome: *letter*, 303
mutation analysis, Crigler-Najjar syndrome type II resulting from three different mutations in the bilirubin uridine 5'-diphosphate-glucuronosyltransferase (*UGT1A1*) gene: *letter*, 712
MPS II in females: molecular basis of two different cases: *letter*, e29
Mutation analysis of *H19* and *NAP1L4* (*hNAP2*) candidate genes and *IGF2* DMR2 in Beckwith-Wiedemann syndrome: *letter*, 212
Mutation analysis of the methyl-CpG binding protein 2 gene (*MECP2*) in patients with Rett syndrome: *letter*, 608
mutation screening, Importance of the autosomal recessive retinitis pigmentosa locus on 1q31-q32.1 (RP12) and mutation analysis of the candidate gene *RGS16* (*RGS-r*): *letter*, 384
Mutation screening in Rett syndrome patients, 250
Screening British CADASIL families for mutations in the *NOTCH3* gene: *letter*, 224
mutation spectrum, The mutation spectrum in Holt-Oram syndrome: *letter*, 785
mutational analysis, Mutational analysis of the human pancreatic secretory trypsin inhibitor (*PSTI*) gene in hereditary and sporadic chronic pancreatitis: *letter*, 67
mutations, Mutations in the *MECP2* gene in a cohort of girls with Rett syndrome: *letter*, 610
***MYOC*,** Novel mutation in the *MYOC* gene in primary open angle glaucoma patients: *letter*, 301
myopia, The heritability of high myopia: a reanalysis of Goldschmidt's data: *letter*, 227
β myosin heavy chain, First description of germline mosaicism in familial hypertrophic cardiomyopathy, 132
β myosin heavy chain mutation, Expression of HCM causing mutations: lessons learnt from genotype-phenotype studies of the South African founder *MYH7* A797T mutation: *letter*, 951

***NAG* amplification,** Gene amplification in PNETs/medulloblastomas: mapping of a novel amplified gene within the *MYCN* amplicon, 501
nail-patella syndrome, Achondroplasia and nail-patella syndrome: the compound phenotype: *letter*, e25
neocentromere, An analphoid supernumerary marker chromosome derived from chromosome 3 ascertained in a fetus with multiple malformations: *letter*, 807
neonatal screening, Many ΔF508 heterozygote neonates with transient hypertrypsinaemia have a second, mild *CFTR* mutation: *letter*, 543
neural tube defects, Comorbidity of 5,10-methylenetetrahydrofolate reductase and methionine synthase gene polymorphisms and risk for neural tube defects: *letter*, 949
neurodevelopmental delay, Deletion (2)(p14p15) in a child with severe neurodevelopmental delay: *letter*, e21
neurofibromatosis, A clinical assessment of neurofibromatosis type 1 (NF1) and segmental NF in northern Finland: *letter*, e43
A search for evidence of somatic mutations in the *NF1* gene, 44
Epidemiology of neurofibromatosis type 1 (NF1) in northern Finland: *letter*, 632
NF2 gene deletion in a family with a mild phenotype: *letter*, 75
neurofibromatosis 1, Growth in North American white children with neurofibromatosis 1 (NF1), 933
neurofibromatosis 2, Clinical and molecular correlates of somatic mosaicism in neurofibromatosis 2: *letter*, 542
neurofibromatosis type 1 (NF1), Tandem duplication within the neurofibromatosis type 1 gene (*NF1*) and reciprocal t(15;16)(q26.3;q12.1) translocation in familial association of NF1 with intestinal neuronal dysplasia type B (IND B): *letter*, 146
neurofibromatosis type 2, Use of MRI and audiological tests in presymptomatic diagnosis of type 2 neurofibromatosis (NF2): *letter*, 944
New Zealand Maori, Genetics of club foot in Maori and Pacific people, 680
NF, A clinical assessment of neurofibromatosis type 1 (NF1) and segmental NF in northern Finland: *letter*, e43
NF1, A clinical assessment of neurofibromatosis type 1 (NF1) and segmental NF in northern Finland: *letter*, e43
***NF1* gene,** A search for evidence of somatic mutations in the *NF1* gene, 44
NF2, *NF2* gene deletion in a family with a mild phenotype: *letter*, 75
Neurofibromatosis type 2: *review*, 897
Niemann-Pick disease type C, Genotype-phenotype relationship of Niemann-Pick disease type C: a possible correlation between clinical onset and levels of NPC1 protein in isolated skin fibrob'asts: *letter*, 707
non-attendance, Why patients do not attend for their appointments at a genetics clinic: *letter*, 810
non-directiveness, Shared decision making and non-directiveness in genetic counselling, 135
non-syndromic hearing loss, Prevalent connexin 26 gene (*GJB2*) mutations in Japanese, 41
Noonan syndrome, Detailed mapping, mutation analysis, and intragenic polymorphism identification in candidate Noonan syndrome genes *MYL2*, *DCN*, *EPS8*, and *RPL6* : *letter*, 884
NOTCH signalling, *JAGGED1* expression in human embryos: correlation with the Alagille syndrome phenotype, 658
***NOTCH3*,** Screening British CADASIL families for mutations in the *NOTCH3* gene: *letter*, 224
Nova Scotia, Absence of fragile X syndrome in Nova Scotia: *letter*, 77
NPC1 protein, Genotype-phenotype relationship of Niemann-Pick disease type C: a possible correlation between clinical onset and levels of NPC1 protein in isolated skin fibroblasts: *letter*, 707
***NRAMP1*,** Identification of novel alleles at a polymorphic microsatellite repeat region in the human *NRAMP1* gene promoter: analysis of allele frequencies in primary biliary cirrhosis: *letter*, 150
nucleotide excision repair, Rough skin, brittle hair, and photosensitivity: a mild phenotypic variant of trichothiodystrophy: *letter*, 312

occipitofrontal circumference, Growth in North American white children with neurofibromatosis 1 (NF1), 933
ocular angioma, Risk of multisystem disease in isolated ocular angioma (haemangioblastoma): *letter*, 62
ocular refraction, The heritability of high myopia: a reanalysis of Goldschmidt's data: *letter*, 227
oculocutaneous albinism type 2, Oculocutaneous albinism type 2 with a *P* gene missense mutation in a patient with Angelman syndrome: *letter*, 392
oesophageal atresia, Interstitial deletion of chromosome 17 (del(17)(q22q23.3 confirms a link with oesophageal atresia: *letter*, 701
ophthalmoplegia, Cardiovascular fibrosis, hydrocephalus, ophthalmoplegia, and visceral involvement in an American child with Gaucher disease: *letter*, e40
osseous abnormalities, Hall-Riggs syndrome: a possible second affected family?: *letter*, 886
osteochondrodysplasia, Anticipation in progressive diaphyseal dysplasia: *letter*, 394
osteogenesis imperfecta, Glycine to tryptophan substitution in type I collagen in a patient with OI type III: a unique collagen mutation, 371

***P* gene missense mutation,** Oculocutaneous albinism type 2 with a *P* gene missense mutation in a patient with Angelman syndrome: *letter*, 392
***p53*,** Absence of germline $p16^{INK4a}$ alterations in *p53* wild type Li-Fraumeni syndrome families: *letter*, e13
pachygyria, New MR/MCA syndrome with distinct facial appearance and general habitus, broad and webbed neck, hypoplastic inverted nipples, epilepsy, and pachygyria of the frontal lobes: *letter*, 460
palaeopathology, A case of dyschondrosteosis from Roman Britain: *letter*, e27
palmoplantar hyperkeratosis, A connexin 26 mutation causes a syndrome of sensorineural hearing loss and palmoplantar hyperkeratosis (MIM 148350), 50
pancreatitis, A CGC>CAT gene conversion-like event resulting in the R122H mutation in the cationic trypsinogen gene and its implication in the genotyping of pancreatitis: *letter*, e36
Papillon-Lefèvre syndrome, Haim-Munk syndrome and Papillon-Lefèvre syndrome are allelic mutations in cathepsin C, 88
Of palms, soles, and gums: *commentary*, 81
parental origin, Parental origin and mechanisms of formation of cytogenetically recognisable de novo direct and inverted duplications, 281
parental origin of alleles, Age and sex based genetic locus heterogeneity in type 1 diabetes, 186
Parkinson's disease, Dipeptidyl carboxypeptidase 1 (*DCP1*) and butyrylcholinesterase (*BCHE*) gene interactions with the apolipoprotein E ε4 allele as risk factors in Alzheimer's disease and in Parkinson's disease with coexisting Alzheimer pathology, 766
partial monosomy 22q, Cryptic subtelomeric translocations in the 22q13 deletion syndrome, 58
partial tetrasomy 3q, Skin pigmentary anomalies in a mosaic form of partial tetrasomy 3q: *letter*, 723
partial trisomy, Partial trisomy 22 in a liveborn resulting from a rearrangement between chromosomes 6 and 22: *letter*, e22
partial trisomy 9q, Molecular cytogenetic characterisation of partial trisomy 9q in a case with pyloric stenosis and a review, 529
partial trisomy/monosomy, An unbalanced submicroscopic translocation t(8;16)(q24.3;p13.3)pat associated with tuberous sclerosis complex, adult polycystic kidney disease, and hypomelanosis of Ito, 287
patella, Breast hypoplasia and disproportionate short stature in the ear, patella, short stature syndrome: expansion of the phenotype?: *letter*, 719
Genitopatellar syndrome: a new condition comprising absent patellae, scrotal hypoplasia, renal anomalies, facial dysmorphism, and mental retardation, 520
***PAX4*,** Exclusion of a disease relevant role of *PAX4* in the aetiology of Silver-Russell syndrome: screening for mutations and determination of imprinting status: *letter*, e44

PCR-DGGE, Analysis of the human tumour necrosis factor-alpha (TNFα) gene promoter polymorphisms in children with bone cancer: *letter,* **789**

PEG1/MEST, Diagnosis of maternal uniparental disomy of chromosome 7 with a methylation specific PCR assay: *letter,* **e19**

periodontal disease, Localisation of a gene for prepubertal periodontitis to chromosome 11q14 and identification of a cathepsin C gene mutation, **95**

peroxisome proliferator activated receptors, Over-representation of *PPARγ* sequence variants in sporadic cases of glioblastoma multiforme: preliminary evidence for common low penetrance modifiers for brain tumour risk in the general population, **410**

phenotype, Genotype-phenotype correlation in three homozygotes and nine compound heterozygotes for the cystic fibrosis mutation 2183AA→G shows a severe phenotype: *letter,* **307**

The genetics of childhood cataract: *review,* **481**

phenotype-genotype correlation, Identification of supernumerary marker chromosomes derived from chromosomes 5, 6, 19, and 20 using FISH, **114**

phenotypic variability, No evidence for imprinting in distal 18q: *letter,* **152**

phosphomannomutase, Identification of four novel *PMM2* mutations in congenital disorders of glycosylation (CDG) Ia French patients, **579**

phosphorylase kinase, Phosphorylase kinase deficient liver glycogenosis: progression to cirrhosis in infancy associated with *PHKG2* mutations (H144Y and L225R): *letter,* **376**

pili torti et canaliculi, Pili torti et canaliculi and agenesis of the teeth: report of a new "pure" hair-tooth ectodermal dysplasia in a Norwegian family: *letter,* **721**

Pitt-Rogers-Danks syndrome, Fine molecular mapping of the 4p16.3 aneuploidy syndromes in four translocation families: *letter,* **449**

PKD1, An unbalanced submicroscopic translocation t(8;16)(q24.3;p13.3)pat associated with tuberous sclerosis complex, adult polycystic kidney disease, and hypomelanosis of Ito, **287**

***PMM2* mutations,** Identification of four novel *PMM2* mutations in congenital disorders of glycosylation (CDG) Ia French patients, **579**

***PMP22* gene,** Hereditary neuropathy with liability to pressure palsies: two cases with a reciprocal translocation t(16;17)(q12;p11.2) interrupting the *PMP22* gene: *letter,* **396**

PNET, Gene amplification in PNETs/medulloblastomas: mapping of a novel amplified gene within the *MYCN* amplicon, **501**

point mutation, Prevalence of mitochondrial gene mutations among hearing impaired patients, **38**

polymorphism, Analysis of the human tumour necrosis factor-alpha (TNFα) gene promoter polymorphisms in children with bone cancer: *letter,* **789**

polymorphisms, *RET* genotypes comprising specific haplotypes of polymorphic variants predispose to isolated Hirschsprung disease, **572**

population based, Novel germline mutations in the *PTEN* tumour suppressor gene found in women with multiple cancers, **336**

port wine stain, A family with hereditary port wine stain: *letter,* **e12**

predictive genetic testing, Evaluation of a counselling protocol for predictive genetic testing for hereditary non-polyposis colorectal cancer, **108**

predictive testing, Attitudes to genetic testing for breast cancer susceptibility in women at increased risk of developing hereditary breast cancer: *letter,* **472**

premature ovarian failure, Autosomal translocation associated with premature ovarian failure: *letter,* **e2**

prenatal diagnosis, Attitudes of von Hippel-Lindau disease patients towards presymptomatic genetic diagnosis in children and prenatal diagnosis: *letter,* **476**

Attitudes towards termination of pregnancy in subjects who underwent presymptomatic testing for the *BRCA1/BRCA2* gene mutation in The Netherlands: *letter,* **883**

Prenatal detection of trisomy for the entire long arm of chromosome 7: *letter,* **551**

prepubertal periodontitis, Localisation of a gene for prepubertal periodontitis to chromosome 11q14 and identification of a cathepsin C gene mutation, **95**

presacral mass, Autosomal dominant sacral agenesis: Currarino syndrome: *review,* **561**

presymptomatic, Use of MRI and audiological tests in presymptomatic diagnosis of type 2 neurofibromatosis (NF2): *letter,* **944**

presymptomatic genetic diagnosis, Attitudes of von Hippel-Lindau disease patients towards presymptomatic genetic diagnosis in children and prenatal diagnosis: *letter,* **476**

presymptomatic testing, Ten years of presymptomatic testing for Huntington's disease: the experience of the UK Huntington's Disease Prediction Consortium, **567**

prevalence, Epidemiology of neurofibromatosis type 1 (NF1) in northern Finland: *letter,* **632**

primary biliary cirrhosis (PBC), Identification of novel alleles at a polymorphic microsatellite repeat region in the human *NRAMP1* gene promoter: analysis of allele frequencies in primary biliary cirrhosis: *letter,* **150**

primary pulmonary hypertension, Sporadic primary pulmonary hypertension is associated with germline mutations of the gene encoding BMPR-II, a receptor member of the TGF-β family, **741**

procollagen processing, Clinical phenotypes and molecular characterisation of three patients with Ehlers-Danlos syndrome type VII: *letter,* **e33**

progressive diaphyseal dysplasia, Anticipation in progressive diaphyseal dysplasia: *letter,* **394**

Localisation of the gene causing diaphyseal dysplasia Camurati-Engelmann to chromosome 19q13, **245**

progressive external ophthalmoplegia, Further evidence for genetic heterogeneity of autosomal dominant disorders with accumulation of multiple deletions of mitochondrial DNA: *letter,* **547**

progressive nephropathy, Constitutional *WT1* mutations correlate with clinical features in children with progressive nephropathy: *letter,* **698**

prostate cancer, No evidence of germline *PTEN* mutations in familial prostate cancer: *letter,* **210**

prostate cancer gene, Suggestive evidence for a site specific prostate cancer gene on chromosome 1p36: *letter,* **947**

protein truncation test, 2157delG: a frequent mutation in *BRCA2* missed by PTT: *letter,* **e42**

protein truncation test (PTT), Diagnostic analysis of Rubinstein-Taybi syndrome: five cosmids should be used for microdeletion detection and low number of protein truncating mutations, **168**

prothrombin, Hemiplegic cerebral palsy and the factor V Leiden mutation: *letter,* **787**

proximal 10q trisomy, Proximal 10q trisomy: a new case with anal atresia: *letter,* **e24**

pseudoachondroplasia, Molecular diagnosis is important to confirm suspected pseudoachondroplasia: *letter,* **64**

pseudoautosomal region, Homozygous deletion of *SHOX* in a mentally retarded male with Langer mesomelic dysplasia: *letter,* **959**

pseudoisodicentric chromosome, Partial tetrasomy 21 in a male infant: *letter,* **e30**

psychological characteristics, Why do women attend familial breast cancer clinics?, **197**

PTEN, Absence of germline mutations in *MINPP1,* a phosphatase encoding gene centromeric of *PTEN,* in patients with Cowden and Bannayan-Riley- Ruvalcaba syndrome without germline *PTEN* mutations: *letter,* **715**

Identification of *PTEN* mutations in metastatic melanoma specimens, **653**

No evidence of germline *PTEN* mutations in familial prostate cancer: *letter,* **210**

Will the real Cowden syndrome please stand up: revised diagnostic criteria: *commentary,* **828**

pulmonary agenesis, Unilateral lobar pulmonary agenesis in sibs: *letter,* **557**

pulmonary lymphangioleiomyomatosis, Mutational analysis of the tuberous sclerosis gene *TSC2* in patients with pulmonary lymphangioleiomyomatosis, **55**

punctate calcification of epiphyses, Punctate calcification of the epiphyses, visceral malformations, and craniofacial dysmorphism in a female baby: *letter,* **796**

pyloric stenosis, Delineation of a new syndrome: clustering of pyloric stenosis, endometriosis, and breast cancer in two families: *letter,* **794**

Molecular cytogenetic characterisation of partial trisomy 9q in a case with pyloric stenosis and a review, **529**

Q576R mutation, No linkage or association of the *IL-4Rα* gene Q576R mutation with atopic asthma in Italian families: *letter,* **382**

questionnaire, The psychological impact of a cancer family history questionnaire completed in general practice: *letter,* **470**

R122H mutation, A CGC>CAT gene conversion-like event resulting in the R122H mutation in the cationic trypsinogen gene and its implication in the genotyping of pancreatitis: *letter,* **e36**

RB1, Two novel germline mutations of the retinoblastoma gene (*RB1*) that show incomplete penetrance, one splice site and one missense: *letter,* **e6**

RBM, Xp;Yp translocation inherited from the father in an *SRY, RBM,* and *TSPY* positive true hermaphrodite with oligozoospermia: *letter,* **e28**

reading disability, Two translocations of chromosome 15q associated with dyslexia, **771**

reasons for attending, Why do women attend familial breast cancer clinics?, **197**

receptor tyrosine kinase, Multiple endocrine neoplasia type 2 and *RET*: from neoplasia to neurogenesis: *review,* **817**

recessive, Mutation analysis of the spastin gene (*SPG4*) in patients with hereditary spastic paraparesis, **759**

recessive inheritance, Unilateral lobar pulmonary agenesis in sibs: *letter,* **557**

reciprocal translocation t(16;17)(q12;p11.2), Hereditary neuropathy with liability to pressure palsies: two cases with a reciprocal translocation t(16;17)(q12;p11.2) interrupting the *PMP22* gene: *letter*, **396**
recurrence risks, Recurrence risks in undiagnosed mental retardation: *letter*, **e45**
recurrent mutation, Recurrent germline mutation in *MSH2* arises frequently de novo, **646**
renal angiomyolipomata, Renal angiomyolipomata and learning difficulty in tuberous sclerosis complex: *letter*, **156**
renal anomalies, Genitopatellar syndrome: a new condition comprising absent patellae, scrotal hypoplasia, renal anomalies, facial dysmorphism, and mental retardation, **520**
renal dysplasia, An HDR (hypoparathyroidism, deafness, renal dysplasia) syndrome locus maps distal to the DiGeorge syndrome region on 10p13/14, **33**
restriction fragment length polymorphism (RFLP), Identification of novel alleles at a polymorphic microsatellite repeat region in the human *NRAMP1* gene promoter: analysis of allele frequencies in primary biliary cirrhosis: *letter*, **150**
retinal dystrophy, Importance of the autosomal recessive retinitis pigmentosa locus on 1q31-q32.1 (RP12) and mutation analysis of the candidate gene *RGS16* (*RGS-r*): *letter*, **384**
retinitis pigmentosa, Genetic heterogeneity of Usher syndrome type II: localisation to chromosome 5q, **256**
Importance of the autosomal recessive retinitis pigmentosa locus on 1q31-q32.1 (RP12) and mutation analysis of the candidate gene *RGS16* (*RGS-r*): *letter*, **384**
Mutation analysis of *GABRR1* and *GABRR2* in autosomal recessive retinitis pigmentosa (*RP25*): *letter*, **e5**
retinoblastoma, A constitutional homozygous mutation in the *RB1* gene in a patient with unilateral retinoblastoma: *letter*, **615**
Two novel germline mutations of the retinoblastoma gene (*RB1*) that show incomplete penetrance, one splice site and one missense: *letter*, **e6**
Rett syndrome, Mutation analysis of the methyl-CpG binding protein 2 gene (*MECP2*) in patients with Rett syndrome: *letter*, **608**
Mutation screening in Rett syndrome patients, **250**
Mutations in the *MECP2* gene in a cohort of girls with Rett syndrome: *letter*, **610**
Spectrum of mutations in the *MECP2* gene in patients with infantile autism and Rett syndrome: *letter*, **e41**
RET, Characterisation of the human *GFRα-3* locus and investigation of the gene in Hirschsprung disease, **674**
Multiple endocrine neoplasia type 2 and *RET*: from neoplasia to neurogenesis: *review*, **817**
revised diagnostic criteria, Will the real Cowden syndrome please stand up: revised diagnostic criteria: *commentary*, **828**
ring chromosome 7, Silver-Russell syndrome and ring chromosome 7: *letter*, **380**
risk assessment, Efficacy of a touchscreen computer based family cancer history questionnaire and subsequent cancer risk assessment, **354**
RLGS, Gene amplification in PNETs/medulloblastomas: mapping of a novel amplified gene within the *MYCN* amplicon, **501**
RP25, Mutation analysis of *GABRR1* and *GABRR2* in autosomal recessive retinitis pigmentosa (*RP25*): *letter*, **e5**
Rubinstein-Taybi syndrome (RTS), Diagnostic analysis of Rubinstein-Taybi syndrome: five cosmids should be used for microdeletion detection and low number of protein truncating mutations, **168**
Rudd syndrome, Interstitial deletion in Xp22.3 is associated with X linked ichthyosis, mental retardation, and epilepsy, **600**
ruptured intracranial aneurysms, The *ACE* I allele is associated with increased risk for ruptured intracranial aneurysms, **498**

sacral agenesis, Autosomal dominant sacral agenesis: Currarino syndrome: *review*, **561**
SALL1, A *SALL1* mutation causes a branchio-oto-renal syndrome-like phenotype: *letter*, **458**
Clinical and molecular cytogenetic studies of a large de novo interstitial deletion 16q11.2-16q21 including the putative transcription factor gene *SALL1* : *letter*, **389**
Three novel *SALL1* mutations extend the mutational spectrum in Townes-Brocks syndrome: *letter*, **303**
Sanfilippo syndrome type A, Mutational analysis of Sanfilippo syndrome type A (MPS IIIA): identification of 13 novel mutations: *letter*, **704**
Sanjad-Sakati syndrome, Confirmation of the assignment of the Sanjad-Sakati (congenital hypoparathyroidism) syndrome (OMIM 241410) locus to chromosome lq42-43: *letter*, **63**
Hypoparathyroidism, retarded growth and development, and dysmorphism or Sanjad-Sakati syndrome: an Arab disease reminiscent of Kenny-Caffey syndrome: *letter*, **145**
sarcoglycans, Disruption of heart sarcoglycan complex and severe cardiomyopathy caused by β sarcoglycan mutations, **102**
sarcolemmal complex proteins, A cross section of autosomal recessive limb-girdle muscular dystrophies in 38 families, **361**
satellited chromosomes, Monosomy and trisomy 1q44-qter in two sisters originating from a half cryptic 1q;15p translocation: *letter*, **612**
scanning electron microscopy, Pili torti et canaliculi and agenesis of the teeth: report of a new "pure" hair-tooth ectodermal dysplasia in a Norwegian family: *letter*, **721**
sclerosing bone dysplasia, Localisation of the gene causing diaphyseal dysplasia Camurati-Engelmann to chromosome 19q13, **245**
screening, Can hair be used to screen for breast cancer?: *letter*, **297**
segmental NF, A clinical assessment of neurofibromatosis type 1 (NF1) and segmental NF in northern Finland: *letter*, **e43**
sensorineural hearing loss, A connexin 26 mutation causes a syndrome of sensorineural hearing loss and palmoplantar hyperkeratosis (MIM 148350), **50**
sequence variants, Over-representation of *PPARγ* sequence variants in sporadic cases of glioblastoma multiforme: preliminary evidence for common low penetrance modifiers for brain tumour risk in the general population, **410**
serotonin, Cockayne syndrome associated with low CSF 5-hydroxyindole acetic acid levels: *letter*, **553**
severe early onset periodontitis, Identification of cathepsin C mutations in ethnically diverse Papillon-Lefèvre syndrome patients, **927**
sex, Age and sex based genetic locus heterogeneity in type 1 diabetes, **186**
sex reversal, Sex reversal and diaphragmatic hernia in phenotypically female sibs with normal XY chromosomes: *letter*, **315**
shared decision making, Shared decision making and non-directiveness in genetic counselling, **135**
SHFM, Delineation of a complex karyotypic rearrangement by microdissection and CGH in a family affected with split foot, **442**
short stature, Clinical and radiographic features of a family with hypochondroplasia owing to a novel Asn540Ser mutation in the fibroblast growth factor receptor 3 gene: *letter*, **220**
short stature homeobox gene, Homozygous deletion of *SHOX* in a mentally retarded male with Langer mesomelic dysplasia: *letter*, **959**
short stature syndrome, Breast hypoplasia and disproportionate short stature in the ear, patella, short stature syndrome: expansion of the phenotype?: *letter*, **719**
sibs, Unilateral lobar pulmonary agenesis in sibs: *letter*, **557**
Silver-Russell syndrome, Biallelic expression of *IGFBP1* and *IGFBP3*, two candidate genes for the Silver-Russell syndrome: *letter*, **65**
Diagnosis of maternal uniparental disomy of chromosome 7 with a methylation specific PCR assay: *letter*, **e19**
Exclusion of a disease relevant role of *PAX4* in the aetiology of Silver-Russell syndrome: screening for mutations and determination of imprinting status: *letter*, **e44**
Silver-Russell syndrome and ring chromosome 7: *letter*, **380**
skeletal disorder, Idiopathic multicentric osteolysis presents early and is not linked to chromosome 18q21.1: *letter*, **e34**
skeletal dysplasia, Clinical and radiographic features of a family with hypochondroplasia owing to a novel Asn540Ser mutation in the fibroblast growth factor receptor 3 gene: *letter*, **220**
Liebenberg syndrome: brachydactyly with joint dysplasia (MIM 186550): a second family: *letter*, **548**
skin, Of palms, soles, and gums: *commentary*, **81**
skin pigmentation, Skin pigmentary anomalies in a mosaic form of partial tetrasomy 3q: *letter*, **723**
***SLC19A2* gene,** A novel mutation in the thiamine responsive megaloblastic anaemia gene *SLC19A2* in a patient with deficiency of respiratory chain complex I, **669**
SLC23A1, Mapping of the human genes (*SLC23A2* and *SLC23A1*) coding for vitamin C transporters 1 and 2 (SVCT1 and SVCT2) to 5q23 and 20p12, respectively: *letter*, **e20**
SLC23A2, Mapping of the human genes (*SLC23A2* and *SLC23A1*) coding for vitamin C transporters 1 and 2 (SVCT1 and SVCT2) to 5q23 and 20p12, respectively: *letter*, **e20**
Slovak population, Novel mutations in the homogentisate-1,2-dioxygenase gene identified in Slovak patients with alkaptonuria: *letter*, **539**
SMAD genes, Mutation analysis of *SMAD2*, *SMAD3*, and *SMAD4* genes in hereditary non-polyposis colorectal cancer: *letter*, **298**
Smith-Lemli-Opitz syndrome, Incidence and molecular mechanism of aberrant splicing owing to a G→C splice acceptor site mutation causing Smith-Lemli-Opitz syndrome: *letter*, **387**
The Smith-Lemli-Opitz syndrome: *review*, **321**
Smith-Magenis syndrome, Circadian rhythm abnormalities of melatonin in Smith-Magenis syndrome, **428**
somatic mosaic, Germline and somatic mosaicism in achondroplasia: *letter*, **956**
somatic mosaicism, Clinical and molecular correlates of somatic mosaicism in neurofibromatosis 2: *letter*, **542**
Somatic mosaicism associated with a mild Alport syndrome phenotype: *letter*, **238**
somatic mutations, A search for evidence of somatic mutations in the *NF1* gene, **44**
spastic paraplegia, A de novo complex chromosomal rearrangement involving chromosomes 2, 3, and 10 associated with microcephaly and early onset spasticity: *letter*, **892**
spastin, Mutation analysis of the spastin gene (*SPG4*) in patients with hereditary spastic paraparesis, **759**

spectral karyotyping, CHARGE association in a child with de novo chromosomal aberration 46,X,der(X)t(X;2)(p22.1;q33) detected by spectral karyotyping: *letter,* **e47**
spermatogenesis, Increase of FMRP expression, raised levels of *FMR1* mRNA, and clonal selection in proliferating cells with unmethylated fragile X repeat expansions: a clue to the sex bias in the transmission of full mutations?, **842**
splice donor site of exon 5, Recurrent germline mutation in *MSH2* arises frequently de novo, **646**
splice site, Three novel *SALL1* mutations extend the mutational spectrum in Townes-Brocks syndrome: *letter,* **303**
splice site mutation, A constitutional homozygous mutation in the *RB1* gene in a patient with unilateral retinoblastoma: *letter,* **615**
Two novel germline mutations of the retinoblastoma gene (*RB1*) that show incomplete penetrance, one splice site and one missense: *letter,* **e6**
spondyloepiphyseal dysplasia congenita, Report of five novel and one recurrent *COL2A1* mutations with analysis of genotype-phenotype correlation in patients with a lethal type II collagen disorder, **263**
SRY, Xp;Yp translocation inherited from the father in an *SRY*, *RBM*, and *TSPY* positive true hermaphrodite with oligozoospermia: *letter,* **e28**
stability, FRAXA and FRAXE: the results of a five year survey, **415**
***STAT6*,** Variants of *STAT6* (signal transducer and activator of transcription 6) in atopic asthma: *letter,* **380**
stature, Growth in North American white children with neurofibromatosis 1 (NF1), **933**
Stickler syndrome, Clinical variability of Stickler syndrome with a *COL2A1* haploinsufficiency mutation: implications for genetic counselling: *letter,* **318**
structural rearrangement mosaicism, The first three mosaic cri du chat syndrome patients with two rearranged cell lines: *letter,* **967**
subtelomeric probes, Perfect endings: a review of subtelomeric probes and their use in clinical diagnosis: *review,* **401**
subtelomeric rearrangement, Cryptic subtelomeric translocations in the 22q13 deletion syndrome, **58**
sulphamidase, Mutational analysis of Sanfilippo syndrome type A (MPS IIIA): identification of 13 novel mutations: *letter,* **704**
supernumerary marker chromosome, A supernumerary marker chromosome originating from two different regions of chromosome 18, **121**
supernumerary marker chromosomes, Identification of supernumerary marker chromosomes derived from chromosomes 5, 6, 19, and 20 using FISH, **114**
survival motor neuron protein, Does the survival motor neuron protein (SMN) interact with Bcl-2?: *letter,* **536**
syndrome, A new X linked mental retardation (XLMR) syndrome with short stature, small testes, muscle wasting, and tremor localises to Xq24-q25, **663**

t(4p;7p), Unexpected high frequency of de novo unbalanced translocations in patients with Wolf-Hirschhorn syndrome (WHS): *letter,* **798**
t(4p;8p), Unexpected high frequency of de novo unbalanced translocations in patients with Wolf-Hirschhorn syndrome (WHS): *letter,* **798**
T705I, I705 variant in the low density lipoprotein receptor gene has no effect on plasma cholesterol levels: *letter,* **713**
tandem duplication, Parental origin and mechanisms of formation of cytogenetically recognisable de novo direct and inverted duplications, **281**
teeth, Of palms, soles, and gums: *commentary,* **81**
telomeres, Perfect endings: a review of subtelomeric probes and their use in clinical diagnosis: *review,* **401**
teratogen, A clinical study of 57 children with fetal anticonvulsant syndromes, **489**
termination of pregnancy, Attitudes towards termination of pregnancy in subjects who underwent presymptomatic testing for the *BRCA1*/*BRCA2* gene mutation in The Netherlands: *letter,* **883**
tetrasomy, Partial tetrasomy 21 in a male infant: *letter,* **e30**
TGF-β, Sporadic primary pulmonary hypertension is associated with germline mutations of the gene encoding BMPR-II, a receptor member of the TGF-β family, **741**
Tietz syndrome, Tietz syndrome (hypopigmentation/deafness) caused by mutation of *MITF* , **446**
TNFα, Analysis of the human tumour necrosis factor-alpha (TNFα) gene promoter polymorphisms in children with bone cancer: *letter,* **789**
tooth agenesis, Pili torti et canaliculi and agenesis of the teeth: report of a new "pure" hair-tooth ectodermal dysplasia in a Norwegian family: *letter,* **721**
Townes-Brocks syndrome, A *SALL1* mutation causes a branchio-oto-renal syndrome-like phenotype: *letter,* **458**
Clinical and molecular cytogenetic studies of a large de novo interstitial deletion 16q11.2-16q21 including the putative transcription factor gene *SALL1* : *letter,* **389**
Three novel *SALL1* mutations extend the mutational spectrum in Townes-Brocks syndrome: *letter,* **303**
transcriptional regulation, Chromatin modification and disease: *review,* **905**
translocation, Autosomal translocation associated with premature ovarian failure: *letter,* **e2**
Two translocations of chromosome 15q associated with dyslexia, **771**
translocations, Disease associated balanced chromosome rearrangements: a resource for large scale genotype-phenotype delineation in man, **858**
transmission disequilibrium test, *RET* genotypes comprising specific haplotypes of polymorphic variants predispose to isolated Hirschsprung disease, **572**
trichothiodystrophy, Rough skin, brittle hair, and photosensitivity: a mild phenotypic variant of trichothiodystrophy: *letter,* **312**
triplet repeat instability, Increase of FMRP expression, raised levels of *FMR1* mRNA, and clonal selection in proliferating cells with unmethylated fragile X repeat expansions: a clue to the sex bias in the transmission of full mutations?, **842**
triplication, Tandem triplication of chromosome 13q14 with inverted interstitial segment in a 4 year old girl: *letter,* **964**
trisomy, Dysmorphic sibs trisomic for the region 6q22.1→6q23.3: *letter,* **889**
trisomy 1q, Monosomy and trisomy 1q44-qter in two sisters originating from a half cryptic 1q;15p translocation: *letter,* **612**
trisomy 20p, Pure trisomy 20p resulting from isochromosome formation and whole arm translocation: *letter,* **454**
trisomy 7q, Prenatal detection of trisomy for the entire long arm of chromosome 7: *letter,* **551**
TRMA syndrome, A novel mutation in the thiamine responsive megaloblastic anaemia gene *SLC19A2* in a patient with deficiency of respiratory chain complex I, **669**
tRNA$^{Ser(UCN)}$, A novel mutation in the mitochondrial tRNA$^{Ser(UCN)}$ gene in a family with non-syndromic sensorineural hearing impairment, **692**
true hermaphroditism, Xp;Yp translocation inherited from the father in an *SRY*, *RBM*, and *TSPY* positive true hermaphrodite with oligozoospermia: *letter,* **e28**
tryptophan, Glycine to tryptophan substitution in type I collagen in a patient with OI type III: a unique collagen mutation, **371**
***TSC2*,** An unbalanced submicroscopic translocation t(8;16)(q24.3;p13.3)pat associated with tuberous sclerosis complex, adult polycystic kidney disease, and hypomelanosis of Ito, **287**
Characterisation of six large deletions in *TSC2* identified using long range PCR suggests diverse mechanisms including Alu mediated recombination: *letter,* **877**
Mutational analysis of the tuberous sclerosis gene *TSC2* in patients with pulmonary lymphangioleiomyomatosis, **55**
TSPY, Xp;Yp translocation inherited from the father in an *SRY*, *RBM*, and *TSPY* positive true hermaphrodite with oligozoospermia: *letter,* **e28**
tuberous sclerosis, Genotype-phenotype correlations in tuberous sclerosis: *letter,* **e2**
tuberous sclerosis complex, Renal angiomyolipomata and learning difficulty in tuberous sclerosis complex: *letter,* **156**
tumour suppressor, Novel germline mutations in the *PTEN* tumour suppressor gene found in women with multiple cancers, **336**
Over-representation of *PPARγ* sequence variants in sporadic cases of glioblastoma multiforme: preliminary evidence for common low penetrance modifiers for brain tumour risk in the general population, **410**
type 1 diabetes, Age and sex based genetic locus heterogeneity in type 1 diabetes, **186**
type II collagen disorders, Report of five novel and one recurrent *COL2A1* mutations with analysis of genotype-phenotype correlation in patients with a lethal type II collagen disorder, **263**
type X collagen, Equal expression of type X collagen mRNA from mutant and wild type *COL10A1* alleles in growth plate cartilage from a patient with metaphyseal chondrodysplasia type Schmid: *letter,* **627**

U type exchange, Tandem triplication of chromosome 13q14 with inverted interstitial segment in a 4 year old girl: *letter,* **964**
undiagnosed mental retardation, Recurrence risks in undiagnosed mental retardation: *letter,* **e45**
universal primer quantitative fluorescent multiplex PCR (UPQFM-PCR), Universal primer quantitative fluorescent multiplex (UPQFM) PCR: a method to detect major and minor rearrangements of the low density lipoprotein receptor gene, **272**
UPD(7)mat, Diagnosis of maternal uniparental disomy of chromosome 7 with a methylation specific PCR assay: *letter,* **e19**
Usher syndrome, Genetic heterogeneity of Usher syndrome type II: localisation to chromosome 5q, **256**

variant, I705 variant in the low density lipoprotein receptor gene has no effect on plasma cholesterol levels: *letter,* **713**
variants, Variants of *STAT6* (signal transducer and activator of transcription 6) in atopic asthma: *letter,* **380**
vascular dementia, Screening British CADASIL families for mutations in the *NOTCH3* gene: *letter,* **224**
vascular malformation, A family with hereditary port wine stain: *letter,* **e12**
vestibular schwannomma, Neurofibromatosis type 2: *review,* **897**

***VHL*,** Cryptic von Hippel-Lindau disease: germline mutations in patients with haemangioblastoma only, **939**
Familial clear cell renal cell carcinoma (FCRC): clinical features and mutation analysis of the *VHL*, *MET*, and *CUL2* candidate genes, **348**
visceral malformations, Punctate calcification of the epiphyses, visceral malformations, and craniofacial dysmorphism in a female baby: *letter*, **796**
vitamin C transporters 1 and 2, Mapping of the human genes (*SLC23A2* and *SLC23A1*) coding for vitamin C transporters 1 and 2 (SVCT1 and SVCT2) to 5q23 and 20p12, respectively: *letter*, **e20**
vocal cord paralysis, Distal spinal muscular atrophy with vocal cord paralysis (dSMA-VII) is not linked to the *MPD2* locus on chromosome 5q31: *letter*, **e14**
von Hippel-Lindau disease, Attitudes of von Hippel-Lindau disease patients towards presymptomatic genetic diagnosis in children and prenatal diagnosis: *letter*, **476**
Cryptic von Hippel-Lindau disease: germline mutations in patients with haemangioblastoma only, **939**

Waardenburg syndrome, Tietz syndrome (hypopigmentation/deafness) caused by mutation of *MITF* , **446**
whistling face, Progressive neurological deterioration in a child with distal arthrogryposis and whistling face: *letter*, **231**
whole arm, Pure trisomy 20p resulting from isochromosome formation and whole arm translocation: *letter*, **454**
Wolf-Hirschhorn syndrome, Fine molecular mapping of the 4p16.3 aneuploidy syndromes in four translocation families: *letter*, **449**
Unexpected high frequency of de novo unbalanced translocations in patients with Wolf-Hirschhorn syndrome (WHS): *letter*, **798**
Wolfram syndrome, The mitochondrial genome in Wolfram syndrome: *letter*, **463**
***WT1* gene,** Sex reversal and diaphragmatic hernia in phenotypically female sibs with normal XY chromosomes: *letter*, **315**
Constitutional *WT1* mutations correlate with clinical features in children with progressive nephropathy: *letter*, **698**
***WT1* mutation,** Familial Wilms tumour resulting from *WT1* mutation: intronic polymorphism causing artefactual constitutional homozygosity: *letter*, **377**

X chromosome, Characterisation and genetic mapping of a new X linked deafness syndrome, **836**
X linked ichthyosis, Interstitial deletion in Xp22.3 is associated with X linked ichthyosis, mental retardation, and epilepsy, **600**
X linked mental retardation, A new X linked mental retardation (XLMR) syndrome with short stature, small testes, muscle wasting, and tremor localises to Xq24-q25, **663**
Interstitial deletion in Xp22.3 is associated with X linked ichthyosis, mental retardation, and epilepsy, **600**
***XNP/ATR-X*,** *ATR-X* mutations cause impaired nuclear location and altered DNA binding properties of the XNP/ATR-X protein, **746**
Xp22.3 deletion, Interstitial deletion in Xp22.3 is associated with X linked ichthyosis, mental retardation, and epilepsy, **600**
Xq24-q25, A new X linked mental retardation (XLMR) syndrome with short stature, small testes, muscle wasting, and tremor localises to Xq24-q25, **663**

Y chromosome, Divergent outcomes of intrachromosomal recombination on the human Y chromosome: male infertility and recurrent polymorphism, **752**
Y chromosome stature gene, FISH deletion mapping defines a single location for the Y chromosome stature gene, *GCY* , **593**

zinc finger, *ATR-X* mutations cause impaired nuclear location and altered DNA binding properties of the XNP/ATR-X protein, **746**